

# **The Engaging Primary Care in Cancer Survivorship (EPICS) study\***

**Protocol Number\* : <Pending>**

**National Clinical Trial (NCT) Identified Number: <Number, once assigned by  
CT.gov>**

**Principal Investigator\*: Erin E. Hahn, PhD, MPH**

**Sponsor: Kaiser Permanente Southern California, Department of Research and  
Evaluation**

**Grant Title: The EPICS (Engaging Primary Care in Cancer Survivorship) study: A  
randomized trial of novel models of care for cancer survivors**

**Grant Number\*: 1R01CA249419-01**

**Funded by: National Cancer Institute (NCI)**

**Version Number: v.1.0**

**January 14, 2021**

### Summary of Changes from Previous Version:

| Affected Section(s) | Summary of Revisions Made                                 | Rationale |
|---------------------|---|-----------|
|                     | Not applicable; this is the first version of the protocol |           |
|                     |   |           |

## **CONFIDENTIALITY STATEMENT**

This document is confidential communication. Acceptance of this document constitutes agreement by the recipient that no unpublished information contained herein will be published or disclosed without prior approval of the Principal Investigator or other participating study leadership and as consistent with the NIH terms of award.

## Table of Contents

|   |    |
|---|----|
| STATEMENT OF COMPLIANCE .....   | 1  |
| INVESTIGATOR'S SIGNATURE.....   | 2  |
| 1     PROTOCOL SUMMARY.....   | 3  |
| 1.1     Synopsis.....   | 3  |
| 1.2     Schema .....  | 4  |
| 1.3     Schedule of Activities .....  | 6  |
| 2     INTRODUCTION .....  | 7  |
| 2.1     Study Rationale.....  | 7  |
| 2.2     Risk/Benefit Assessment.....  | 9  |
| 2.2.1     Known Potential Risks.....  | 9  |
| 2.2.2     Known Potential Benefits .....  | 10 |
| 2.2.3     Assessment of Potential Risks and Benefits.....   | 10 |
| 3     OBJECTIVES AND ENDPOINTS .....  | 11 |
| 4     STUDY DESIGN.....   | 12 |
| 4.1     Overall Design.....   | 12 |
| 4.2     Scientific Rationale for Study Design.....  | 14 |
| 4.3     Justification for Intervention .....  | 14 |
| 4.4     End-of-Study Definition .....   | 15 |
| 5     STUDY POPULATION .....  | 15 |
| 5.1     Inclusion Criteria .....  | 15 |
| 5.2     Exclusion Criteria .....  | 16 |
| 5.3     Lifestyle Considerations.....   | 16 |
| 5.4     Screen Failures.....  | 16 |
| 5.5     Strategies for Recruitment and Retention.....   | 17 |
| 6     STUDY INTERVENTION(S) OR EXPERIMENTAL MANIPULATION(S).....  | 19 |
| 6.1     Study Intervention(s) or Experimental Manipulation(s) Administration.....                                     | 19 |
| 6.1.1     Study Intervention or Experimental Manipulation Description.....  | 19 |
| 6.1.2     Administration and/or Dosing .....  | 20 |
| 6.2     Fidelity .....  | 21 |
| 6.2.1     Interventionist Training and Tracking .....   | 21 |
| 6.3     Measures to Minimize Bias: Randomization and Blinding.....  | 21 |
| 6.4     Study Intervention/Experimental Manipulation Adherence.....   | 22 |
| 6.5     Concomitant Therapy .....   | 22 |
| 6.5.1     Rescue Therapy .....  | 22 |
| 7     STUDY INTERVENTION/EXPERIMENTAL MANIPULATION DISCONTINUATION AND<br>PARTICIPANT DISCONTINUATION/WITHDRAWAL..... | 22 |
| 7.1     Discontinuation of Study Intervention/Experimental Manipulation .....   | 22 |
| 7.2     Participant Discontinuation/Withdrawal from the Study .....   | 22 |
| 7.3     Lost to Follow-Up .....   | 23 |
| 8     STUDY ASSESSMENTS AND PROCEDURES .....  | 23 |
| 8.1     Endpoint and Other Non-Safety Assessments.....  | 23 |
| 8.2     Safety Assessments.....   | 25 |
| 8.3     Adverse Events and Serious Adverse Events.....  | 26 |
| 8.3.1     Definition of Adverse Events .....  | 26 |
| 8.3.2     Definition of Serious Adverse Events.....   | 26 |

|         |   |    |
|---------|---|----|
| 8.3.3   | Classification of an Adverse Event.....                           | 26 |
| 8.3.4   | Time Period and Frequency for Event Assessment and Follow-Up..... | 27 |
| 8.3.5   | Adverse Event Reporting.....                                      | 28 |
| 8.3.6   | Serious Adverse Event Reporting .....                             | 28 |
| 8.3.7   | Reporting Events to Participants .....                            | 28 |
| 8.3.8   | Events of Special Interest .....                                  | 29 |
| 8.3.9   | Reporting of Pregnancy .....                                      | 29 |
| 8.4     | Unanticipated Problems.....                                       | 29 |
| 8.4.1   | Definition of Unanticipated Problems .....                        | 29 |
| 8.4.2   | Unanticipated Problems Reporting.....                             | 29 |
| 8.4.3   | Reporting Unanticipated Problems to Participants .....            | 30 |
| 9       | STATISTICAL CONSIDERATIONS .....                                  | 30 |
| 9.1     | Statistical Hypotheses.....                                       | 30 |
| 9.2     | Sample Size Determination.....                                    | 30 |
| 9.3     | Populations for Analyses .....                                    | 32 |
| 9.4     | Statistical Analyses.....   | 32 |
| 9.4.1   | General Approach.....   | 32 |
| 9.4.2   | Analysis of the Primary Endpoint(s) .....                         | 33 |
| 9.4.3   | Analysis of the Secondary Endpoint(s).....                        | 34 |
| 9.4.4   | Safety Analyses.....  | 34 |
| 9.4.5   | Baseline Descriptive Statistics .....                             | 34 |
| 9.4.6   | Planned Interim Analyses .....                                    | 34 |
| 9.4.7   | Sub-Group Analyses .....  | 34 |
| 9.4.8   | Tabulation of Individual Participant Data .....                   | 35 |
| 9.4.9   | Exploratory Analyses.....   | 35 |
| 10      | SUPPORTING DOCUMENTATION AND OPERATIONAL CONSIDERATIONS .....     | 35 |
| 10.1    | Regulatory, Ethical, and Study Oversight Considerations.....      | 35 |
| 10.1.1  | Informed Consent Process .....                                    | 35 |
| 10.1.2  | Study Discontinuation and Closure .....                           | 36 |
| 10.1.3  | Confidentiality and Privacy .....                                 | 36 |
| 10.1.4  | Future Use of Stored Specimens and Data .....                     | 37 |
| 10.1.5  | Key Roles and Study Governance .....                              | 38 |
| 10.1.6  | Safety Oversight.....   | 38 |
| 10.1.7  | Clinical Monitoring.....  | 38 |
| 10.1.8  | Quality Assurance and Quality Control.....                        | 39 |
| 10.1.9  | Data Handling and Record Keeping.....                             | 40 |
| 10.1.10 | Protocol Deviations.....  | 40 |
| 10.1.11 | Publication and Data Sharing Policy .....                         | 40 |
| 10.1.12 | Conflict of Interest Policy .....                                 | 43 |
| 10.2    | Additional Considerations.....                                    | 43 |
| 10.3    | Abbreviations and Special Terms .....                             | 43 |
| 10.4    | Protocol Amendment History .....                                  | 45 |
| 11      | REFERENCES .....  | 46 |

## STATEMENT OF COMPLIANCE

The trial will be carried out in accordance with International Council on Harmonisation Good Clinical Practice (ICH GCP) and the following:

- United States (US) Code of Federal Regulations (CFR) applicable to clinical studies (45 CFR Part 46, 21 CFR Part 50, 21 CFR Part 56, 21 CFR Part 312, and/or 21 CFR Part 812).

National Institutes of Health (NIH)-funded investigators and clinical trial site staff who are responsible for the conduct, management, or oversight of NIH-funded clinical trials have completed Human Subjects Protection and ICH GCP Training.

The protocol, informed consent form(s), recruitment materials, and all participant materials will be submitted to the IRB for review and approval. Approval of both the protocol and the consent form(s) must be obtained before any participant is consented. Any amendment to the protocol will require review and approval by the IRB before the changes are implemented to the study. All changes to the consent form(s) will be IRB approved; a determination will be made regarding whether a new consent needs to be obtained from participants who provided consent, using a previously approved consent form.

## INVESTIGATOR'S SIGNATURE

The signature below constitutes the approval of this protocol and provides the necessary assurances that this study will be conducted according to all stipulations of the protocol, including all statements regarding confidentiality, and according to local legal and regulatory requirements and applicable US federal regulations and ICH guidelines.

Principal Investigator or Clinical Site Investigator:

Signed:

Date:

\_\_\_\_\_  
Name<sup>\*</sup> : Erin E. Hahn, PhD, MPH

Title<sup>\*</sup> : Research Scientist / Associate Professor

### Investigator Contact Information

Affiliation<sup>\*</sup> : Kaiser Permanente Southern California, Department of Research and Evaluation  
Kaiser Permanente Bernard J. Tyson School of Medicine, Department of Health Systems Science  
Address: 100 S. Los Robles, 2<sup>nd</sup> Floor, Pasadena, CA 91101  
Telephone: 626-564-3505  
Email: Erin.E.Hahn@kp.org

## 1 PROTOCOL SUMMARY


### 1.1 SYNOPSIS

|                           |  |
|---------------------------|--|
| <b>Title:</b>             | The Engaging Primary Care in Cancer Survivorship (EPICS) study   |
| <b>Grant Number:</b>      | 1R01CA249419-01  |
| <b>Study Description:</b> | <p>A quasi-experimental non-randomized pre/post with control group trial of two models of cancer survivorship care in early-stage colorectal and breast cancer survivors cared for in a community-based, integrated health care setting. The trial will test the efficacy of an embedded primary care provider (PCP) model (experimental condition) in which PCPs are embedded within an oncology practice and will care for low-risk survivors who will be transitioned at 6-12 months post-treatment for comprehensive survivorship care. We hypothesize that a) patients in the PCP model will have superior receipt of recommended care compared to usual care; b) patients in the PCP model will perceive significantly better care coordination, self-efficacy, and confidence in their PCP compared to usual care; and c) use of unplanned and non-recommended care will be less in the PCP model compared to usual care.</p> |
| <b>Objectives* :</b>      | <p>Primary Objective: To determine the efficacy of an embedded PCP model (experimental condition) compared to usual care on use of recommended cancer surveillance and preventive care services.</p> <p>Secondary Objectives: 1) To determine the efficacy of an embedded PCP model (experimental condition) compared to usual care on patient-reported outcomes; 2) to compare use of unplanned and non-recommended care in the PCP model compared to usual care.</p>   |
| <b>Endpoints* :</b>       | <p>Primary Endpoint: Receipt of guideline-recommended cancer surveillance and preventive care services assessed over a 36-month period.</p> <p>Secondary Endpoints: 1) Patient reported outcomes: Validated measures of patient and provider communication, and coordination of survivorship care; important covariates will include quality of life, physical and mental health, assessment of survivorship care delivery, and satisfaction with health decisions; 2) Receipt of non-recommended surveillance care based on current clinical guidelines, and use of non-planned hospitalization, emergency department, and urgent care services.</p>  |



|   |  |
|---|--|
| <b>Study Population:</b>  | Patients: 3,750 adult (21+) Kaiser Permanente Southern California members diagnosed and treated for first primary early-stage breast (stage 0, I, II) or colorectal (stage I, II) cancer within Kaiser Permanente Southern California (breast cancer patients may be receiving trastuzumab or endocrine therapy). Patients will be at low-risk for recurrence and treatment-related toxicities, as determined by our risk algorithm. Providers: For centers assigned to the embedded PCP model, PCPs selected to participate must be Board Certified in a relevant primary care specialty; hold a valid and current MD or advanced practitioner license; and be employed by the Southern California Permanente Medical Group.  |
| <b>Phase * or Stage:</b>  | This is a single site phase III National Institutes of Health Clinical Trial   |
| <b>Description of Sites/Facilities Enrolling Participants:</b>      | This pre/post with control group trial will be conducted in a community-based integrated system, Kaiser Permanente Southern California (KPSC), with medical centers assigned to the embedded PCP or usual care model.  |
| <b>Description of Study Intervention/Experimental Manipulation:</b> | <p>Embedded PCPs will provide comprehensive care for survivors, including cancer surveillance services, preventive care, and management of long-term therapy and associated side effects (e.g., endocrine therapy in breast survivors). A comprehensive multilevel approach will prepare survivors and PCPs (physicians and/or advance practice providers):</p> <p><u>Intervention: Embedded PCP.</u> Embedded PCPs will be enrolled in a 4-month course of initial training and education, followed by ongoing education via tailored survivorship information for low-risk survivors transitioning to their care. Initial training features 3 core components to build capacity, skill, and knowledge: 1) Individual didactic learning; 2) Small in-person group sessions; and 3) Observation.</p> <p><u>Intervention: Patient-level.</u> Eligible patients in the embedded PCP model will be provided with tailored education regarding the planned transition prior to the transition. After cessation of active treatment, the care team will provide printed information including the planned course of survivorship care, what to expect from embedded PCP care, when the transition will occur, and reassurance that the oncology team will be available via telephone and email, and that PCPs will refer back to the oncologist for any concerning signs or symptoms.</p> <p><u>Intervention: System-level.</u> Tailored alerts in the EMR for recommended cancer surveillance and preventive care services that include rationale, links to guidelines, and references for questions (scheduled to fire when survivors have an outpatient office visit with their embedded PCP).</p> |
| <b>Study Duration*:</b>   | 60-months  |
| <b>Participant Duration:</b>  | 36-months  |

## 1.2 SCHEMA



### 1.3 SCHEDULE OF ACTIVITIES

|   | Year 1 |    |    |    | Year 2 |    |    |    | Year 3 |    |    |    | Year 4 |    |    |    | Year 5 |    |    |    |
|---|--------|----|----|----|--------|----|----|----|--------|----|----|----|--------|----|----|----|--------|----|----|----|
|   | Q1     | Q2 | Q3 | Q4 | Q1     | Q2 | Q3 | Q4 | Q1     | Q2 | Q3 | Q4 | Q1     | Q2 | Q3 | Q4 | Q1     | Q2 | Q3 | Q4 |
| Administrative and regulatory tasks   | x      | x  |    |    |        |    |    |    |        |    |    |    |        |    |    |    |        |    |    |    |
| Center assignment   |        | x  | x  |    |        |    |    |    |        |    |    |    |        |    |    |    |        |    |    |    |
| Pre-period data collection and analysis   |        |    | x  | x  | x      | x  |    |    |        |    |    |    |        |    |    |    |        |    |    |    |
| Initiate primary care training at intervention sites                                |        |    | x  | x  |        |    |    |    |        |    |    |    |        |    |    |    |        |    |    |    |
| Annual PCP survey of knowledge, confidence in survivorship care                     |        |    |    |    | x      |    |    |    | x      |    |    |    | x      |    |    |    | x      |    |    |    |
| <b>Patient identification</b> , risk stratification at all sites                    |        |    | x  | x  | x      | x  | x  |    |        |    |    |    |        |    |    |    |        |    |    |    |
| <b>Enrollment period</b> into intervention or usual care, N=3750 (~900 per quarter) |        |    |    | x  | x      | x  | x  | x  |        |    |    |    |        |    |    |    |        |    |    |    |
| <b>Follow-up period</b> for intervention or usual care (36 months)                  |        |    |    |    | x      | x  | x  | x  | x      | x  | x  | x  | x      | x  | x  | x  |        |    |    |    |
| Patient survey at 12 months post-transition (rolling recruitment)                   |        |    |    |    |        |    |    | x  | x      | x  | x  | x  |        |    |    |    |        |    |    |    |
| Survey analyses (Aim 2)   |        |    |    |    |        |    |    |    |        |    |    |    | x      | x  | x  |    |        |    |    |    |
| Refine EMR-based measures of utilization  |        |    |    |    |        |    |    |    |        |    |    |    |        | x  | x  |    |        |    |    |    |
| Obtain EMR data on utilization, initiate analyses (Aims 1, 3)                       |        |    |    |    |        |    |    |    |        |    |    |    |        |    |    | x  | x      | x  |    |    |

## 2 INTRODUCTION

### 2.1 STUDY RATIONALE

**Advances in treatment and detection of cancer have led to a rapidly growing population of cancer survivors.** Significant growth of cancer patients, particularly those  $\geq 65$  years, is predicted within the U.S. over the next 20 years.<sup>1</sup> The majority of these patients will enter into a prolonged post-treatment phase of care and considered cancer survivors.<sup>2</sup> *Cancer survivor* is commonly defined as from the time of diagnosis through the balance of his or her life, as originally articulated by the National Coalition for Cancer Survivorship.<sup>3</sup> For this proposal, we are focusing on the transition from the end of active cancer treatment (e.g., surgery, chemotherapy, radiation) to the post-treatment survivorship phase, which may include extended hormonal or other chronic therapy. Survivors require coordinated, comprehensive care addressing the major domains of survivorship as described by National Academy of Medicine, the American Society for Clinical Oncology (ASCO) and others, including surveillance for recurrence, screening for new cancers, screening and management for long-term and late effects, symptom management, and preventive care (e.g., vaccinations, screening for other diseases).<sup>4-13</sup> Older survivors ( $>65$  years), who make up the majority of cancer survivors, also have a higher prevalence of comorbid conditions requiring management and coordination of care.<sup>1</sup>

**We are facing unprecedented challenges in survivorship care delivery.** Currently, survivorship care delivery is unsystematic, occurs in a variety of settings, and is often poorly coordinated.<sup>14-16</sup> Serious gaps in survivorship care have been identified, including underuse of guideline-recommended cancer surveillance services, such as annual mammography for breast cancer survivors and colonoscopy for colorectal cancer (CRC) survivors,<sup>17-22</sup> and underuse of recommended preventive care services such as vaccinations, lipid testing and cardiovascular risk management, and cancer screenings.<sup>23,24,25-27</sup> In addition, the frequent overuse of non-recommended services that provide limited benefit exposes patients to significant harms, as exemplified by use of non-recommended serum tumor marker tests and high-intensity imaging for surveillance of early stage CRC and breast cancer.<sup>17,28-36</sup> Breast and CRC survivors have characterized their survivorship care as lacking in information for follow-up and providing inadequate support to address their needs.<sup>37</sup> Compounding these issues, our current oncology workforce is insufficient to care for the rapidly growing population of survivors.<sup>1,38,39</sup> Estimates from ASCO show a projected growth in demand of approximately 50% and only 14% growth in clinician supply, with demand exceeding supply as soon as 2020.<sup>40,41</sup> This demand is increasing wait times for oncology services across the U.S.<sup>42</sup> The current approach to delivering survivorship care, which is typically with oncology specialists with limited or no coordination with primary care, is inadequate and fails to meet the needs of many survivors.<sup>6</sup> In this study, we will evaluate a new approach to survivorship care that embeds survivorship-trained primary care providers (PCPs) into the oncology setting to provide survivorship care for low-risk survivors, potentially alleviating demand for oncology services while maintaining quality care.

**PCP-led survivorship care is feasible, but critical barriers must be addressed.** In observational studies, survivors who receive care that involves PCPs (whether physicians or advance practice providers) received higher quality of care.<sup>23,25,43-46</sup> Most survivors expect PCPs to manage their general preventive care.<sup>47</sup> PCPs

are well-positioned to counsel survivors regarding lifestyle choices such as physical activity,<sup>48</sup> disease prevention,<sup>49</sup> screening for psychosocial health needs,<sup>50</sup> and tobacco cessation.<sup>51</sup> Receipt of CVD preventive care in cancer survivors was strongly associated with PCP involvement.<sup>52</sup> PCPs receive high ratings on care coordination and comprehensive care during survivorship.<sup>53</sup> However, lack of knowledge and other barriers must be addressed for PCPs to assume primary responsibility for survivorship care; for example, a recent survey of PCPs found that less than half of guideline-recommended services for breast cancer survivors were routinely used.<sup>54</sup> Additionally, survivors may not be comfortable transitioning to a PCP for survivorship.<sup>55,56</sup> Some survivors may prefer oncology-led cancer surveillance based on emotional reasons and feeling connected to their oncology team.<sup>57,58</sup> Other barriers perceived by clinicians and survivors include poor communication with oncology,<sup>59</sup> lack of knowledge regarding existing guidelines,<sup>60</sup> limitations of health information technology and support,<sup>61</sup> and inadequate survivorship training and education.<sup>58,62,60,63,64,65,66</sup> Survivorship care plans (SCPs), summative documents intended to help improve communication and coordination of survivorship care,<sup>7</sup> may be a necessary but insufficient tool to address these issues. SCPs have been shown to help PCPs feel more prepared and knowledgeable about survivorship care delivery,<sup>67</sup> improve PCP-reported care coordination and communication,<sup>68</sup> and may improve quality of survivorship care<sup>69-71</sup> and patient self-efficacy.<sup>72</sup> However, results from most randomized controlled trials (RCTs) have found little to no effect of SCPs on distress, quality of life, or satisfaction with care.<sup>73,74</sup> Thus, SCPs alone may be a necessary but insufficient component to engage PCPs and improve quality of survivorship care.<sup>73,75-77</sup> We will address these barriers identified in the empirical literature in our study with multi-level interventions (patient, provider, system) incorporating robust education, training, and support for survivorship care.

**The need for risk-stratified survivorship care.** Risk-stratified survivorship care has been suggested as a potential model for U.S. survivors.<sup>10,41,78-80</sup> Risk-stratification involves evaluation of cancer and treatment characteristics, risk of complications, comorbid conditions, and other patient characteristics. In this approach, survivors with less complicated cancer treatments and at low-risk for recurrence or treatment toxicities are triaged to a less intense pathway (e.g., PCP-led or self-management) and patients with more complex cancer treatments/stage of disease are triaged to specialist oncology care for ongoing follow-up.<sup>41,80</sup> This type of stratification has been implemented within the National Health Service in the U.K.,<sup>81-83</sup> and a similar approach has been implemented in Northern Ireland.<sup>84</sup> Recent data show successful triage of half of CRC survivors and approximately 80% of breast survivors to a low-intensity survivorship pathway.<sup>82</sup> In the U.K. the stratified pathways have been found to improve access to oncology specialists and are projected to save approximately £90 million over 5 years.<sup>80</sup> We will integrate a risk-stratified approach in our study to identify and triage low-risk CRC and breast cancer survivors, incorporating these evidence-based recommendations.

To address these issues, we will conduct a non-randomized pre/post with control trial of two models of cancer survivorship care in early-stage colorectal and breast cancer survivors cared for in a community-based, integrated health care setting. Building on our pilot work<sup>57</sup> and drawing on the empirical literature, we will test the efficacy of an embedded PCP model (experimental condition) in which PCPs are embedded within an oncology practice, and will care for low-risk survivors who will be transitioned at 12 months post-treatment for comprehensive survivorship care. These PCPs will be given training in cancer

survivorship care focused on cancer surveillance and chronic disease prevention. This model will be compared to the extant oncology physician model (usual care) that is the default condition in the health care system. The primary outcome will be receipt of guideline-recommended care (e.g., cancer surveillance, preventive care), assessed over a 36-month period. Secondary outcomes will include validated patient-reported outcomes (PROs), as well as utilization of unplanned care (hospitalizations, urgent care) and receipt of non-recommended care. We will also explore PCP perceptions of confidence in delivery of survivorship care. We will conduct this trial in large medical centers of an integrated health care system, Kaiser Permanente Southern California (KPSC), with approximately 3,750 survivors. Data from multiple studies conducted by our team demonstrate the need to improve survivorship care in this setting<sup>31,32</sup> and our recent pilot study demonstrates the feasibility of the embedded PCP model in this setting.<sup>57</sup> KPSC is an exceptional setting to conduct the proposed research with unparalleled access to longitudinal data, a large and diverse patient population, and engaged clinicians.

## 2.2 RISK/BENEFIT ASSESSMENT

### 2.2.1 KNOWN POTENTIAL RISKS

There are minimal physical, psychological, social, legal, privacy or other risks from participating in this study. Individual patients and clinicians will not be recruited for Aims 1 or 3, thus there is minimal risk to loss of privacy or confidentiality; demographic, clinical and utilization data for these Aims will be collected from EMR and/or claims data with no participant interaction.

For this study, a waiver for informed consent will be required to ensure a scientifically valid comparison of the embedded PCP (ePCP) model to current usual care.

We do not foresee any risks to potential participants by not directly consenting them to participate in the trial. This study poses minimal risk to the patient because survivorship cancer care will be performed as part of routine medical care, and both the ePCP and the oncology-based model for survivorship cancer care are consistent with guidelines from professional societies and organizations with regard to the delivery of safe, appropriate and usual care practices. We also believe that such a waiver of consent is not only widely recognized as necessary and appropriate, but also scientifically necessary and ethically justifiable under the “Common Rule” for protection of research participants (45 CFR 46.116d) – (e.g., 1) the research involves no more than minimal risk to the subjects; 2) the waiver of alteration will not adversely affect the rights or welfare of the subjects; 3) the research could not practicably be carried out without the waiver or alteration; and 4) whenever appropriate, the subjects will be provided with additional pertinent information after participation in the form of a summary sheet of survey findings after completion of the study).

Throughout the study phases, we will work closely with our Institutional Review Board, clinical chiefs, and operational leaders to ensure patient safety, with regular review and reporting of adverse events and protocol deviations.

Patients will be individually recruited for patient-reported outcome surveys under Aim 2, initially through email and subsequently through mail and telephone. If email addresses are unavailable, we will use mailed letters as the initial contact. Recruitment scripts will describe the purpose of the research, the selection process for participation, the study structure and time commitment (one-time survey), and risks of participating. We will obtain written or oral consent from all participants for participation in the surveys. Survey consents and opt out options will be built into an electronic template as part of the initial survey screens.

For utilization data captured under Aim 3 from the EMR, individual patients will not be recruited and the ethical justification of the waiver of individual consent, the “Common Rule” for protection of research participants (45 CFR 46.116d), is outlined above.

A rare but serious harm that could occur is loss of confidentiality of personal health information (PHI), but every effort will be made to protect PHI. Specifically, each survey participant will be given a unique alpha-numeric identification number. No data collection form will be linked to an actual participant name. Any document linking the participant name to an identification number will be kept in a locked file separate from data collection files. All records from the study will be kept in locked files. Identifying information will be removed from the final data sets and stored in a linked file to which only the Principal Investigator or designee has access. Individual participants will not be identified in any reports. Additionally, study reports will be aggregated so that individual participants are not identified. All investigators and research staff at participating sites will be required to maintain up-to-date training in human subject protection and good clinical practice through Collaborative Institutional Training Initiative (CITI; <https://www.citiprogram.org/default.asp>) and HIPAA training. Additional security precautions include encryption, digital certification, audit logs, and firewall protection.

---

#### 2.2.2 KNOWN POTENTIAL BENEFITS

There are no direct benefits to the participants in this proposed study. There are potential societal benefits from our findings. Our study is innovative, among the first trials in the U.S. comparing two survivorship models of care and one of the first to use a novel risk-stratified approach that will triage low-risk breast cancer and CRC survivors to one of these models of care. In addition, the study uses a multi-level approach that includes patient-, provider- and system-level components. Furthermore, the ePCP training materials and resources that will comprise a study “toolkit” are mostly free, widely-accessible materials and our risk-stratification methods are easily replicable, making our findings from this study well-poised for scale-up and dissemination.

---

#### 2.2.3 ASSESSMENT OF POTENTIAL RISKS AND BENEFITS

There are minimal physical, psychological, social, legal, privacy or other risks from participating in this study. Individual patients and clinicians will not be recruited for Aims 1 or 3, thus there is minimal risk to loss of privacy or confidentiality; demographic, clinical and utilization data for these Aims will be collected from the EMR and/or claims data with no participant interaction. A rare but serious harm that could occur

is loss of confidentiality of PHI. However, participant confidentiality will be maintained at all times using the specific steps and procedures outlines in section 2.31 above.

This research is essential to advance the field of survivorship cancer care. Definitive evidence on effective models of care that engage primary care physicians and patients in survivorship with equivalent or superior outcomes to oncology-led models would allow clinicians, health system leaders, and policy makers to widely and confidently implement a new approach to survivorship that optimizes patient outcomes and efficiently employs resources.

### 3 OBJECTIVES AND ENDPOINTS

| OBJECTIVES   | ENDPOINTS  | JUSTIFICATION FOR ENDPOINTS  | PUTATIVE MECHANISMS OF ACTION |
|--|--|--|-------------------------------|
| <b>Primary</b>   |  |  |                               |
| To determine the efficacy of an ePCP model (experimental condition) compared to usual care on use of recommended cancer surveillance and preventive care services. | Receipt of guideline-recommended cancer surveillance and preventive care services assessed over a 36-month period  | ePCPs trained to provide care to cancer survivors will provide superior care for preventive services and equivalent cancer surveillance care compared to the usual care model. |                               |
| <b>Secondary</b>   |  |  |                               |
| To determine the efficacy of an ePCP model (experimental condition) compared to usual care on patient-reported outcomes.   | Validated, reliable measures of patient-provider communication, and coordination of survivorship care.<br><br>We will include quality of life, physical and mental health, assessment of survivorship care delivery, and satisfaction with health decisions as important covariates. | ePCPs trained to provide care to cancer survivors will have superior communication and care coordination compared to usual care.   |                               |



| OBJECTIVES  | ENDPOINTS  | JUSTIFICATION FOR ENDPOINTS  | PUTATIVE MECHANISMS OF ACTION |
|---|--|--|-------------------------------|
|   |  |  |                               |
| To determine the efficacy of an ePCP model (experimental condition) compared to usual care on utilization of emergency, urgent, hospital, and non-recommended care. | Utilization: Unplanned care (hospitalizations, emergency department, and urgent care); and receipt of non-recommended cancer surveillance care as described by clinical practice guidelines (e.g., ASCO Choosing Wisely) | ePCPs trained to provide care to cancer survivors may have different rates of patients with unplanned care or non-recommended care.  |                               |
| <b>Tertiary/Exploratory</b>   |  |  |                               |
| To explore knowledge and confidence in survivorship care between ePCPs and PCPs at usual care centers.  | Confidence in delivery of survivorship care and knowledge of survivorship care guidelines, assessed using validated survey questions.  | Causal mechanism: ePCPs exposed to our multi-level intervention of education, training, and support will demonstrate superior knowledge and confidence in delivery of survivorship care compared to PCPs at the usual care centers, who are not exposed. |                               |

## 4 STUDY DESIGN

### 4.1 OVERALL DESIGN

This is a single site (KPSC), multi-center (multiple KPSC medical centers) trial using a non-randomized pre/post control group design to compare the effectiveness of 2 risk-stratified models of survivorship care, an ePCP model with a survivorship-trained PCP providing cancer surveillance, symptom management, and preventive care, compared to an oncology-led model, where the survivor stays with their oncology team (usual care within the KPSC system). We will address the barriers to PCP involvement from the empirical literature and propose to measure recommended quality outcomes including receipt of recommended care, PROs, and utilization of hospital, urgent, and non-recommended care.

This trial will evaluate the efficacy of 2 models of cancer survivorship care on receipt of recommended care for early-stage, low-risk colorectal and breast cancer survivors (primary outcome), PROs (secondary outcome), and use of unplanned and non-recommended care (secondary outcomes). This trial will be conducted in a community-based integrated system, KPSC, with medical centers non-randomly assigned to the ePCP or usual care model, with the models running for 36 months. We will retrospectively (pre-period) and prospectively (post-period) identify early-stage breast and colorectal cancer patients at all sites during their initial cancer treatment using a combination of pathology data and text information extraction algorithms. Potentially eligible survivors will be assigned a binary risk score (low risk/non-low risk) based on data readily available in the EMR including cancer stage, risk of recurrence, and other factors derived from current consensus-based recommendations. Embedded PCP participants will be transitioned to a trained PCP at 6-12 months after the cessation of active treatment. Multi-level components for this model include empirically-derived survivorship care interventions, such as provider education and training, using established training curriculum for PCPs engaging in survivorship care, health IT support (e.g., proactive office encounter alerts, links to guidelines), and patient education on the transition in care, signs and symptoms to be aware of, and what to expect from their PCP for surveillance and prevention. Usual care participants will be followed by their treating oncologist or surgeon, which is current “usual care” within KPSC.

Our specific aims and hypotheses (H) are:

**Aim 1.** To determine the efficacy of the PCP model relative to the usual care control group on receipt of recommended care (primary outcome). We will construct a composite measure of guideline-recommended cancer surveillance and age-appropriate preventive care services for the 18-month pre-period and 18-month post-period, identifying utilization from EMR data.

H1. Patients in the PCP model will have superior receipt of recommended care compared to usual care.

**Aim 2.** To compare patient-centered outcomes on coordination of survivorship care, self-efficacy in managing care, and confidence in their PCP to provide survivorship care between the 2 models (secondary outcomes).

H2. Patients in the PCP model will perceive significantly better care coordination, self-efficacy, and confidence in their PCP compared to usual care

**Aim 3.** To assess utilization of unplanned hospitalizations, use of urgent care, and receipt of non-recommended cancer surveillance services between the 2 models (secondary outcomes).

H3. Use of unplanned and non-recommended care will be significantly less in the PCP model compared to usual care.

**Exploratory Aim.** We will explore PCP knowledge of and confidence in providing survivorship care, comparing PCPs who participate in the embedded PCP model to PCPs practicing at centers assigned to the usual care model.

**Assignment and consent.** Assignment of centers will occur prior to PCP educational interventions and enrollment of patients. A scientifically valid comparison of the embedded PCP model to current usual care would only be possible with population-based enrollment and outcome assessment of low-risk survivors.

Consequently, a waiver of the usual requirement for informed consent will be required. We believe that such a waiver of consent is not only widely recognized as necessary and appropriate but is also scientifically necessary and ethically justified given the minimal risk of the proposed research based on the Common Rule, 45 CFR 46.116d (e.g., use of routine care practices, no investigational drugs or devices, little risk of physical or psychological harm). We will work closely with our Institutional Review Board (IRB), clinical chiefs, and operational leaders to ensure patient safety, with regular review and reporting of adverse events and protocol deviations.

## 4.2 SCIENTIFIC RATIONALE FOR STUDY DESIGN

Despite widespread recognition and interest in survivorship in the U.S., there has been little research on models of survivorship care, or uptake of PCP survivorship models. Extant research has focused mainly on observational studies, such as the recent Patient-Centered Outcomes Research Institute (PCORI)-funded study “Evaluating different types of survivorship care” (study HSRP20143084, Mead et al.) that focused primarily on identifying model types among 236 Commission on Cancer-accredited organizations; a patient survey evaluated PROs. Findings indicate 3 major models of care (specialist consultative, mid-level provider longitudinal, and oncology-based) and few differences in PROs.<sup>85,86</sup> While valuable, the study was not prospective, did not randomize, and did not evaluate important outcomes such as utilization. Eight ‘Survivorship Centers of Excellence’ were developed across the U.S. from 2006-11 with investment from the Lance Armstrong Foundation (one led by Dr. Patricia Ganz in collaboration with Dr. Hahn, Co-I and PI respectively for this proposal). These Centers implemented a variety of different models of care within their cancer centers, but did not conduct head-to-head comparisons of the models as a part of their implementation and research agenda.<sup>87,88</sup> Multiple review papers and an AHRQ technical brief have categorized observed survivorship models of care and developed recommendations on potential model types, content, and structure.<sup>2,6,10,14,15,78,79,89-91</sup> A review by Howell et al. determined that further research is needed to evaluate the efficacy of models of care;<sup>15</sup> McCabe et al. had a similar conclusion.<sup>90</sup> This paucity of high-quality research, particularly in community-based settings, seriously limits our ability to implement policy or practice decisions regarding survivorship care delivery, or to address the impact of clinician shortages and rising health care costs impacting systems, patients, and families.

## 4.3 JUSTIFICATION FOR INTERVENTION

Our proposed research builds on the foundational evidence from observational studies summarized above. Survivorship care provided by oncology specialists alone focuses on cancer surveillance and screening, but often lacks attention to preventive care or management of comorbid conditions. In contrast, care provided by PCPs is more likely to include age appropriate preventive care, as well as management of comorbid conditions and psychosocial concerns, without sacrificing timely detection of cancer recurrence or other adverse events, and may provide increased capacity and access to oncology services for newly-diagnosed patients.<sup>25,43-46</sup> In our pilot study we found an embedded PCP model to be highly feasible and acceptable for low-risk patients.<sup>57</sup> Reviews of survivorship models have identified examples of primary care physician-led or “embedded PCP” models in the U.S., albeit a limited number.<sup>79</sup>

These examples describe setting, domains of care, and care processes (e.g., strategy for managing long-term and late effects).<sup>92,93</sup> However, as noted above, there are few, if any, evaluations of these models.

Collectively, this evidence demonstrates the feasibility and capability of PCPs assuming the primary role in providing survivorship care for early-stage, low-risk survivors, as well as the need for evaluation of survivorship care models. Thus, we will conduct a non-randomized pre/post with control trial to compare the effectiveness of an embedded PCP model, with a survivorship-trained PCP embedded into the oncology clinic to provide comprehensive survivorship care (e.g., cancer surveillance, symptom management, preventive care) as compared to an oncology-led model (usual care), where the survivor stays with their treating oncology team and sees their regular PCP in parallel, but in an uncoordinated fashion. We will address the barriers to PCP involvement identified in the empirical literature in the development of the intervention. Guided by the Framework for Quality Survivorship Care,<sup>11</sup> we propose to measure highly relevant outcomes including receipt of recommended care, PROs, including patient-provider communication and quality of life, and utilization of hospital, urgent, and non-recommended care.

#### 4.4 END-OF-STUDY DEFINITION

A participant is considered to have completed the study if he or she has:

- Been identified and enrolled into either the ePCP or the oncology model of care
- Completed the 36-month follow-up period after enrollment into the intervention
- Completed the survey at 12-months post-transition (rolling recruitment)

### 5 STUDY POPULATION

#### 5.1 INCLUSION CRITERIA

In order to be eligible to participate in this study, an individual must meet all of the following criteria:

**Provider population:**

PCPs selected to participate must be Board Certified in a relevant specialty; hold a valid and current MD or advanced practitioner license; and be employed by the Southern California Permanente Medical Group (SCPMG).

**Patient population:**

- Adult (21+) KPSC members diagnosed and treated for first primary early-stage breast (stage 0, I, II) or colorectal (stage I, II) cancer within KPSC; breast cancer patients may be receiving trastuzumab or endocrine therapy
- At low-risk for recurrence and treatment-related toxicities, as determined by our risk algorithm

For our Aim 2 survey:

- Primary language of English or Spanish (although we will assess our Aim 1 participants for recorded preferred spoken language and adjust translations as needed)
- Ability to complete surveys of patient-reported outcomes

Individuals of all races, ethnicities, and genders are eligible for this study. Children will not be included in the study, because they are rarely, if ever, affected by breast or colorectal cancer.

## 5.2 EXCLUSION CRITERIA

An individual who meets any of the following criteria will be excluded from participation in this study:

- Pre-existing cancer diagnosis (other than non-melanoma skin cancer)
- Less than 120 days of KPSC membership after diagnosis of primary breast cancer or CRC
- Serious medical or psychiatric condition that would detract from study participation and measurement of PROs

For the provider survey, there are no exclusions; all sex/gender, racial, and ethnic group members are eligible to participate.

## 5.3 LIFESTYLE CONSIDERATIONS

N/A

## 5.4 SCREEN FAILURES

We will retrospectively (pre-period) and prospectively (post-period) identify early-stage (stage 0-II) breast and colorectal patients at our assigned intervention condition medical centers early in their treatment course using our comprehensive pathology database, Co-Path, combined with a text-information extraction system. We will identify the end of active treatment (surgery, chemotherapy, and/or radiation) using our treatment databases (e.g., Beacon chemotherapy, Mosaiq radiation) and EMR data. At the end of treatment, we will assess salient risk factors based on recent recommendations for stratifying low-risk survivors. We will create a risk-score (low risk/non-low-risk) based on: stage of disease, risk of recurrence, treatment-related toxicity, and pre-existing comorbidity burden, with assessment of potential interaction of treatment and comorbidity (e.g., breast patients with CVD who receive anthracyclines might be at high-risk for treatment-related CVD events). For patients enrolled in the intervention (post-period), we will send treating oncologists monthly lists of survivors to confirm as low-risk and eligible for the study; treating oncologists may have information beyond what is available in our structured EMR data that may affect risk status. Eligible survivors receiving care in embedded PCP model will receive an automated referral and patient education.

Screen failures are defined as participants with early-stage breast or colorectal cancer who are categorized as low-risk for recurrence and treatment toxicities but are subsequently determined to be not at low risk by either the treating oncologist or ePCP. Patients determined to be not at low-risk will be excluded from the study and will receive their survivorship care under usual care conditions, with their treating oncologist.

## 5.5 STRATEGIES FOR RECRUITMENT AND RETENTION

For Aim 1, there is no individual participant recruitment or contact and patients will not be individually enrolled in this study; rather entire medical centers will be intervention or control with patients passively enrolled. We will prospectively identify early-stage (stage 0-II) breast and colorectal patients at our medical centers assigned to the intervention condition early in their treatment course to introduce the program and initiate tracking. Therefore, anyone who meets the eligibility criteria will be included in either the ePCP or usual care model. Our target enrollment size is 3,750 subjects.

There are no specific strategies that will be used to recruit and retain historically under-represented populations in order to target sample size and conform with the *NIH Policy on Inclusion of Women and Minorities and Inclusion of Individuals Across the Lifespan as Participants in Research Involving Human Subjects*. Individuals of all races, ethnicities, and genders are eligible for this study. KPSC has an extremely diverse member population and no survivor would be excluded from participation in the care models based on spoken language (comprehensive interpreter services are available at all centers). KPSC patients are socioeconomically diverse with insurance coverage that includes commercial plans, Medicare Advantage, Medicaid, and individual insurance with options for high-, medium-, and low-deductible plans, and reside in urban, suburban, and rural communities. KPSC medical center characteristics are similar on distribution of age and gender, with distribution of race/ethnicity echoing the race/ethnicity distribution of Southern California.

Children will not be included in the study, because they are rarely, if ever, affected by breast or colorectal cancer.

Estimated distribution of age for early-stage breast and colorectal cancer survivors:

|                 | <b>Breast</b> | <b>Colorectal</b> |
|-----------------|---------------|-------------------|
| <b>Mean Age</b> | 61.2          | 65.3              |
| <b>&lt;=40</b>  | 5%            | 2%                |
| <b>41-50</b>    | 16%           | 11%               |
| <b>51-60</b>    | 26%           | 23%               |
| <b>61-70</b>    | 31%           | 28%               |
| <b>71-80</b>    | 17%           | 23%               |
| <b>81+</b>      | 6%            | 13%               |

Identification of participants will begin in the 4<sup>th</sup> quarter of Year 1.

Under Aim 2, we will use Dillman's Tailored Design Method to guide all survey activities including recruitment. This approach has three fundamental goals: 1) minimize all possible sources of survey error, including coverage, sampling, nonresponse, and measurement error. Our mixed mode approach will keep our sources of error low by reducing survey burden and increasing coverage; 2) tailor to the survey population by developing survey and recruitment methods that consider participant characteristics and that interact and work together to encourage response; 3) use procedures that create positive social exchange and encourage response. We designed our recruitment procedures to encourage response and minimize nonresponse by using multiple modes of contact, multiple modes of response (e.g., direct link to web survey, paper survey), and incentives. Mixed-mode strategies such as the one we propose have been shown to be effective and efficient and help minimize total survey error. Key Tailored Design features of our approach include: sending a pre-incentive and promising completion incentives, using unified design elements across modes, using multiple modes of communication and multiple contacts, and using recruitment materials tailored for the study population (e.g., stressing the knowledge gained to improve cancer survivorship care).

We will send an introductory letter with a \$2 pre-completion incentive to all potential participants. Participants will receive a \$20 gift card for completing the survey. These are nominal incentive levels widely used in current studies within our department for survey completion; therefore, we anticipate no undue influence or coercion by providing this level of incentive to potential participants.

The invitation letter will include a short URL for the web survey leading directly to the survey and will include a toll-free number and email for opting out of further contact or asking questions. We will use KPSC branding to provide legitimacy for KPSC members. The letter will stress the potential impact of knowledge gained from the survey in helping future cancer survivors and the importance of the participant's expertise in this topic, following social exchange theory principles. We will also indicate our intention to share the results with participants to further establish trust and improve response. We will include a self-addressed stamped return envelope for paper surveys. Non-responders will be sequentially contacted first by email (when available), mail (with a paper survey included), and finally by telephone (for an interviewer-administered survey).

A secure internal study database will permit us to track the recruitment status of each member of the original survey sample at every phase. We will conduct rolling recruitment over a 1-year period by making ordered contacts in waves. Using real-time reporting, we will continually assess response rates and adjust the timing and order of contacts, as needed, to most effectively achieve the desired sample size and diversity (i.e. sex, race/ethnicity). The content of recruitment materials may also be modified based on participants' feedback and response. By employing multiple avenues for recruitment (mail, phone, and email), we can ensure a strong response.

For Aim 3, there is no participant contact; we will obtain utilization data from the KPSC electronic medical record and associated research databases for eligible participants at the embedded PCP (experimental) and usual care arm. This aim is to determine utilization of unplanned hospitalizations, urgent care, and

non-recommended cancer surveillance services, using CPT and ICD codes to identify hospitalizations related to cancer, use of urgent care, and use of non-recommended care. Non-recommended care measures will be identified from relevant guidelines (e.g., ASCO Choosing Wisely, American Cancer Society (ACS) Survivorship).

**Exploratory Aim.** We will invite our ePCPs and PCPs from our usual care arm to participate in a brief, online survey of knowledge and confidence in cancer survivorship care. Recruitment methods for KPSC clinicians will include email, letter, and/or telephone outreach. The overall PCP population at KPSC (N=2,932) consists of 34% white, 49% Asian, 11% Hispanic, and 5% black; 48% are female. We anticipate our PCP sample at participating centers to reflect these characteristics.

#### Retention.

Aims 1 and 3 will rely on a passive retention approach, as there is not participant contact and patients will not be individually enrolled in this study. Patients who maintain active KPSC membership status will be retained.

Under Aim 2, retention is not applicable, because the PRO survey is a one-time cross-sectional study; we will invite all eligible survivors to participate in the PRO survey at 12 months after their transition to the ePCP or the oncology model. Therefore, we do not propose active follow-up of participants who participate in this one-time survey.

## 6 STUDY INTERVENTION(S) OR EXPERIMENTAL MANIPULATION(S)

### 6.1 STUDY INTERVENTION(S) OR EXPERIMENTAL MANIPULATION(S) ADMINISTRATION

#### 6.1.1 STUDY INTERVENTION OR EXPERIMENTAL MANIPULATION DESCRIPTION

Embedded PCPs will provide comprehensive care for survivors, including cancer surveillance services, preventive care, and management of long-term therapy and associated side effects (e.g., endocrine therapy in breast survivors). We have designed a comprehensive multilevel approach to prepare survivors and PCPs (physicians and/or advance practice providers).

**Intervention: Embedded PCP.** We will enroll ePCPs in a 4-month course of initial training and education, followed by ongoing education via tailored survivorship information for low-risk survivors transitioning to their care. We will use widely and freely available resources to help ensure generalizability and scalability of the training approach. Initial training features 3 core components to build capacity, skill, and knowledge, guided by the ASCO Core Curriculum for Survivorship Training: 1) Individual didactic learning; 2) Small in-person group sessions; and 3) Observation. For individual learning, we will use the “Cancer Survivorship E-Learning Series for Primary Care Providers” training modules, a freely available online educational series developed by expert clinicians, researchers, and survivors at the National Cancer Survivorship Resource Center in collaboration with the ACS. Next, small group sessions with the research team, including clinical co-investigators from primary care and oncology, to review survivorship resources



and address questions for 60 minutes each session for 4 weeks, with pre-session work of 1-2 hours per week for sessions 1-3. Lastly, observational learning will take place via shadowing oncologists for 5-10 hours per week for 2 weeks, again leveraging PCP education time, and attending tumor boards (ongoing). We will conduct a brief annual online survey to assess PCP knowledge and confidence in survivorship care and will compare ePCPs to PCPs in the usual care arm (Exploratory Aim). When survivors transition to the ePCP, we will include a tailored packet of educational information and reminders for the ePCP: the ACS survivorship guidelines for breast or CRC cancer, which are designed to assist PCPs in care of survivors,. We will include a standardized letter reminding the ePCP about goals of care and how to work with the oncology team. Importantly, this letter will include when to refer the survivor back to oncology/surgery. Typically, this will include unremitting symptoms such as shortness of breath or pain.

**Intervention: Patient-level.** Eligible patients in the ePCP model will be provided with tailored education regarding the planned transition prior to the transition, a critical factor according to our formative evaluation. After cessation of active treatment, the care team will provide printed information including the planned course of survivorship care, what to expect from ePCP care, when the transition will occur, and reassurance that the oncology team will be available via telephone and email, and that PCPs will refer back to the oncologist for any concerning signs or symptoms. This will be included as part of the standard KPSC survivorship binder provided at the end of treatment, as well as uploaded to the kp.org patient portal.

**Intervention: System-level.** Working with our health IT team, we will create tailored alerts in the EMR for recommended cancer surveillance and preventive care services that include rationale, links to guidelines, and references for questions. These will fire when survivors have an outpatient office visit with their ePCP.

## 6.1.2 ADMINISTRATION AND/OR DOSING

**Intervention: Embedded PCP.** We will engage ePCPs in a 4-month course of initial training and education, followed by ongoing education via tailored survivorship information for low-risk survivors transitioning to their care. Initial training features 3 core components to build capacity, skill, and knowledge: 1) Individual didactic learning; 2) Small in-person group sessions; and 3) Observation.

| Embedded PCP survivorship training activities |   |                                  |
|---|---|----------------------------------|
| Weeks 1-8                                     | Didactic learning from online survivorship care for PCPs modules <sup>94</sup>  |                                  |
| Module 1, 2                                   | Overview; role of PCPs  | Late effects, symptom management |
| Module 3, 4                                   | Psychosocial needs  | Preventive care                  |
| Module 5, 6                                   | Care coordination   | Cancer recovery                  |
| Module 7, 8                                   | Colorectal survivors  | Breast survivors                 |
| Weeks 9-12                                    | Small group sessions with research team and clinical leaders  |                                  |
| Session 1                                     | <u>Pre-session work:</u> View videos: 1) <i>Long-Term Survivorship Care After Cancer Treatment</i> <sup>95</sup> ; 2) <i>From Cancer Patient to Cancer Survivor: Lost in Transition</i> <sup>96</sup><br><u>In session:</u> 1) Review organization and expectations of care. 2) Review and discuss information, resources from <i>Cancer Survivorship in Primary Care</i> <sup>97</sup> |                                  |
| Session 2                                     | <u>Pre-session work:</u> Review <i>ASCO Treatment Plan and Summary Resources</i> <sup>98</sup>  |                                  |

|                 |   |
|-----------------|---|
|                 | <u>In session</u> : Review ASCO Survivorship Compendium; live examples of KPSC Epic-based SCP; review how to use and update SCP in Epic   |
| Session 3       | <u>Pre-session work</u> : Review <i>ACS Survivorship Guidelines</i> for breast, CRC<br><u>In-session work</u> : Discuss ACS guidelines; review planned health-IT support                        |
| Session 4       | <u>In-session work</u> : Review embedded PCP care protocol, when to refer back to oncology, and other workflow processes (e.g., identification and referral process), Q&A with clinical leaders |
| Weeks 13-16     | Observation   |
| Weeks 13-14     | Shadow breast and colorectal oncologists to observe cancer care processes   |
| Weeks 13-16+    | Attend tumor boards and oncology staff meetings   |
| Care transition | Tailored information including: the survivor's SCP with link to the EMR, the breast or colorectal ACS guideline, standardized letter with goals of care   |
| Evaluation      | Brief online survey on perceived confidence and knowledge of survivorship care; <sup>60</sup> evaluation of training content  |

When survivors transition to ePCP care, we will include a tailored packet of educational information: the ACS survivorship guidelines for breast or CRC cancer, which are designed to assist PCPs in care of survivors, a copy and link to the survivor's SCP housed within the EMR (routinely completed by the oncology teams using the Epic template). We will include a standardized letter reminding the embedded PCP about goals of care and how to work with the oncology team. Importantly, this letter will include when to refer the survivor back to oncology/surgery. Typically, this will include unremitting symptoms such as shortness of breath or pain.

**Intervention: Patients.** After cessation of active treatment, the care team will provide printed information including the planned course of survivorship care, what to expect from ePCP care, when the transition will occur, and reassurance that the oncology team will be available via telephone and email, and that PCPs will refer back to the oncologist for any concerning signs or symptoms. This will be included as part of the standard KPSC survivorship binder provided at the end of treatment, as well as uploaded to the patient's kp.org portal.

**Intervention: System-level.** These will fire when survivors have an outpatient office visit with their embedded PCP.

## 6.2 FIDELITY

### 6.2.1 INTERVENTIONIST TRAINING AND TRACKING

We will assess center compliance with the study protocol monthly, with review of relevant data. We will collect utilization data for services with the ePCP, the treating oncology team, and with the patient's assigned KPSC PCP. We will check for patterns of utilization indicative non-compliance with the ePCP protocol (e.g., no or limited visits with the ePCP, continued use of oncology for patients in the ePCP arm). We will have regular interactions with the study sites to discuss and review these data.

## 6.3 MEASURES TO MINIMIZE BIAS: RANDOMIZATION AND BLINDING

N/A. This is quasi-experimental pre/post with control group design.

#### 6.4 STUDY INTERVENTION/EXPERIMENTAL MANIPULATION ADHERENCE

Participant adherence at the patient level does not apply to this study. Participants may or may not have visits with their ePCP and/or oncology team.

#### 6.5 CONCOMITANT THERAPY

N/A

##### 6.5.1 RESCUE THERAPY

N/A

### 7 STUDY INTERVENTION/EXPERIMENTAL MANIPULATION DISCONTINUATION AND PARTICIPANT DISCONTINUATION/WITHDRAWAL

#### 7.1 DISCONTINUATION OF STUDY INTERVENTION/EXPERIMENTAL MANIPULATION

When a subject discontinues from the ePCP model but not from the study, remaining study procedures will be completed as indicated by the study protocol. If a clinically significant finding is identified after enrollment, the investigator or qualified designee will determine if any change in participant management is needed. Any new clinically relevant finding will be reported as an adverse event (AE).

The data to be collected at the time of study intervention discontinuation will include the following:

- The reason(s) for discontinuing the participant from the intervention, and methods for determining the need to discontinue
- Where the participant will be receiving survivorship care.

#### 7.2 PARTICIPANT DISCONTINUATION/WITHDRAWAL FROM THE STUDY

Participants are free to withdraw from participation in the study at any time upon request. An investigator may discontinue a participant from the study for the following reasons:

- Significant study intervention non-compliance, unless varying compliance is an aspect of the study objectives;
- Lost-to-follow up; unable to contact subject (see **Section 7.3, Lost to Follow-Up**)
- Any event or medical condition or situation occurs such that continued collection of follow-up study data would not be in the best interest of the participant or might require an additional treatment that would confound the interpretation of the study;
- The participant meets an exclusion criterion (either newly developed or not previously recognized) that precludes further study participation.

The reason for participant discontinuation or withdrawal from the study will be recorded on the EPICS Case Report Form (CRF). Subjects who receive the study intervention, and subsequently withdraw, or are discontinued from the study, will not be replaced.

### 7.3 LOST TO FOLLOW-UP

For our utilization aims (Aim 1, 3) we will track all utilization within KPSC until disenrollment or death.

For the participant survey, a participant will be considered lost to follow-up if he or she does not respond to multiple contacts to complete the survey after agreeing to do so.

The following actions must be taken if a participant fails to return the survey:

- The site will attempt to contact the participant, resend the survey, and ascertain if the participant wishes to and/or should continue in the study
- Before a participant is deemed lost to follow-up, the investigator or designee will make every effort to regain contact with the participant (where possible, 3 telephone calls and, if necessary, a certified letter to the participant's last known mailing address or local equivalent methods). These contact attempts will be documented in the participant's study file.
- Should the participant continue to be unreachable, he or she will be considered to have withdrawn from the study aim with a primary reason of lost to follow-up.

## 8 STUDY ASSESSMENTS AND PROCEDURES

### 8.1 ENDPOINT AND OTHER NON-SAFETY ASSESSMENTS

**Background Characteristics and Potential Confounders.** Demographic information (e.g., age, race, ethnicity, census-level education and income) and clinical information (e.g., date of cancer diagnosis, age at diagnosis, disease stage, cancer treatment, endocrine therapy) will be collected from the EMR for all participants. Self-reported demographics will be collected from those who participate in the PRO survey. We will collect for clinician variables such as clinician gender, race, and years in practice from the EMR. Sex as a potentially important biologic factor will be included in all analyses.

**Receipt of Recommended Care (Aim 1).** We will construct a composite measure of recommended care for cancer surveillance and for preventive care, based on recommendations from the ACS survivorship guideline. Cancer surveillance for early-stage breast cancer includes annual mammography and 2-4 physician visits with history and physical (H&P) for the first 3 years; CRC surveillance includes colonoscopy within first year, annual chest/abdominal/pelvic CT, and 2-4 H&P visits and CEA lab tests annually for the first 3 years. Preventive care measures will include appropriate cancer screening (other than for surveillance of primary cancer) for CRC, cervical, breast, and lung cancers, vaccinations (flu, pneumococcal), counseling for smoking cessation, and management of hyperlipidemia and

hypercholesterolemia. These recommendations will be tailored to individual age, cancer type, health behaviors, and prior care (e.g., for breast cancer survivors  $\geq 50$ , colonoscopy every 10 years or annual FIT Kit for CRC cancer screening). We will use validated algorithms from our prior work based on CPT and ICD codes to identify use of services.<sup>17,31,32,99-102</sup> We will identify receipt of recommended care for up to 36 months, starting 12 months after cessation of active treatment. Recommended care will be assessed in 12-month periods, with a binary outcome of received/did not receive for recommended cancer surveillance and for preventive care. We will also compare rates of individual services (e.g., rates of vaccination). Survivors with a pattern indicative of recurrence, based on validated recurrence algorithms,<sup>103,104</sup> will be excluded from analysis of recommended cancer care. We will identify use of services from oncology, ePCP, and the member's assigned KPSC PCP to evaluate the impact of the model on utilization of services.

**Patient Reported Outcomes (Aim 2).** We will use validated, reliable measures from existing national surveys that have been used in our target population. Most measures come from well-established surveys such as the Medical Expenditure Panel Survey (MEPS), the European Organisation for the Research and Treatment of Cancer (EORTC) Quality of Life, the Consumer Assessment of Healthcare Providers and Systems (CAHPS), and the Patient-Reported Outcomes Measurement Information System (PROMIS). We will assess patient-reported coordination of care, patient-provider communication, physical and mental health, assessment of survivorship care delivery, satisfaction with health decisions, and other measures relevant to survivorship care. We will also collect socioeconomic, demographic, and health behavior data (e.g. smoking). To evaluate our secondary outcome of patient-provider communication and coordination, we will use validated measures from the MEPS Experiences with Cancer Supplement CAHPS surveys (i.e., Patient-Centered Medical Home CAHPS coordination items). The MEPS patient-provider communication scale evaluates discussions about: 1) cancer follow-up care; 2) late or long-term treatment effects; 3) lifestyle recommendations; and 4) emotional or social needs, with response options discussed in detail, briefly discussed, did not discuss, and I don't remember. We will also examine CAHPS care coordination that includes six items on how often providers had appropriate information about: care provided by specialists; medical records and information about care; patient prescriptions and medications; follow-up regarding patient labs, X-rays, or other tests; and managing care between different providers. The response set for these items is a 4-point Likert scale ranging from "Never" to "Sometimes" to "Usually" to "Always." We will use design principles of Dillman et al. to order questions to minimize testing order effects, ensure the survey is appropriately and consistently designed for multiple modes, and minimize participant burden.<sup>105</sup>

**Utilization of unplanned hospitalizations, urgent care, and non-recommended cancer surveillance services (Aim 3).** Using a similar approach to Aim 1, we will use CPT and ICD codes to identify hospitalizations related to cancer, using primary ICD code to identify reason for admission, as well as use of urgent care and use of non-recommended care. Non-recommended care measures are identified from relevant guidelines (e.g., ASCO Choosing Wisely, ACS Survivorship).<sup>115-117</sup> For early-stage breast cancer, this includes serum tumor markers and high-intensity imaging (e.g., CT and PET scans); for CRC, PET scans.

**Exploratory Aim.** We will explore PCP knowledge of and confidence in providing survivorship care, comparing the embedded PCPs to PCPs practicing at centers assigned to the usual care model.

| <b>Measures for Key Constructs for Patient Survey</b><br>All measures have been previously validated and used in cancer research |   |
|--|---|
| <b>Construct</b>   | <b>Measure</b>  |
| <b>Patient-provider communication about survivorship care</b>  | Medical Expenditure Panel Survey (MEPS) 2016, Experiences with Cancer Supplement, Section 7, Medical Care for Cancer <sup>106</sup> |
| <b>Quality of life</b>   | European Organisation for the Research and Treatment of Cancer Quality of Life Questionnaire Core 30 (EORTC) <sup>107</sup>         |
| <b>Coordination and continuity of cancer care</b>  | Cancer Consumer Assessment of Healthcare Providers and Systems (CAHPS) <sup>108</sup>   |
| <b>Self-efficacy</b>   | PROMIS self-efficacy for managing symptoms <sup>109</sup>   |
| <b>Overall health</b>  | PROMIS Global Health Scale <sup>109</sup>   |
| <b>Receipt of treatment summary and/or care plan</b>   | Receipt of treatment summary from Health Information and National Trends Survey (HINTS 4, Cycle 2) <sup>110,111</sup>               |
| <b>Delivery of survivorship care</b>   | Delivery of Survivorship Care Scale <sup>63</sup>   |
| <b>Symptom burden</b>  | Patient-Reported Bother From Side Effects of Cancer Therapy <sup>112</sup>  |
| <b>Satisfaction with health care decisions</b>   | Patient satisfaction with health care decisions: the satisfaction with decision scale <sup>113</sup>                                |
| <b>Confidence in care</b>  | Questions adapted from 'Confidence in managing survivorship care' Casillas et al <sup>114</sup>                                     |

## 8.2 SAFETY ASSESSMENTS

Collection of observation data from administrative and clinical databases, self-reported demographics from the PRO surveys, and intervention activities at the KPSC medical centers will be collected and discussed at weekly project meetings. During monthly investigator meetings, data and recruitment summary reports of demographic and clinical information will be presented. Recruitment strategies will be brainstormed for optimization as needed. Recruitment success by racial/ethnic categories will be monitored closely and reported in National Institutes of Health (NIH) progress reports and to the Project Officer (PO)/Data Safety Monitoring Board (DSMB) no less than annually.

## 8.3 ADVERSE EVENTS AND SERIOUS ADVERSE EVENTS

### 8.3.1 DEFINITION OF ADVERSE EVENTS

We believe this study poses no more than minimal risk to subjects and the protocol uses the definition of adverse event (AE) from the KPSC IRB: (a) any unfavorable medical or psychological events experienced by a study participant during clinical research, including: a) a new symptom; b) worsening of an existing condition; or a clinically significant abnormal lab finding. AEs are considered a reportable unanticipated problem (UP) -- e.g., reportable to the KPSC IRB -- if they meet all three criteria for an unanticipated problem:

- Unexpected in nature, severity, or frequency
- Related or possibly related to participation in the research
- Suggests greater risk to participants or others than previously known

In the event that an AE is determined, it must be reported to the IRB within 10 business days and at the continuing review period.

### 8.3.2 DEFINITION OF SERIOUS ADVERSE EVENTS

As defined by the KPSC IRB, a serious adverse event (SAE) includes:

- Death
- Life threatening condition/situation
- An enduring or significant incapacity or substantial disruption of the ability to conduct normal life functions
- The delivery of a child with congenital anomaly or birth defect
- Other medical events that the PI determines require intervention to prevent the above outcomes

### 8.3.3 CLASSIFICATION OF AN ADVERSE EVENT

#### 8.3.3.1 SEVERITY OF EVENT

The Principal Investigator must take action to protect the study participant(s) or others from the unexpected risk of harm. For AEs not included in the protocol defined grading system, the following guidelines will be used to describe severity:

- **Mild** - Events require minimal or no treatment and do not interfere with the participant's daily activities.

- **Moderate** – Events result in a low level of inconvenience or concern with the intervention measures. Moderate events may cause some interference with functioning.
- **Severe** – Events interrupt a participant’s usual daily activity and may require systemic drug therapy or other treatment. Severe events are usually potentially life-threatening or incapacitating. Of note, the term “severe” does not necessarily equate to “serious”

---

#### 8.3.3.2 RELATIONSHIP TO STUDY INTERVENTION/EXPERIMENTAL MANIPULATION

All AEs will have their relationship to study procedures, including the intervention, assessed by an appropriately-trained clinician based on temporal relationship and his/her clinical judgment. The degree of certainty about causality will be graded using the categories below.

- **Related** – The AE is known to occur with the study procedures, there is a reasonable possibility that the study procedures caused the AE, or there is a temporal relationship between the study procedures and the event. Reasonable possibility means that there is evidence to suggest a causal relationship between the study procedures and the AE.
- **Not Related** – There is not a reasonable possibility that the study procedures caused the event, there is no temporal relationship between the study procedures and event onset, or an alternate etiology has been established.

---

#### 8.3.3.3 EXPECTEDNESS

The Principal Investigator will determine if the event or incident is unexpected in nature, severity, or frequency, taking into consideration:

- The protocol-related documents, including: 1) IRB-approved research protocol or research application; or 2) other sources of information
- The Principal Investigator’s knowledge of the characteristics of the study population
- The expected progression of any underlying diseases or conditions of the participants
- The participant’s pre-existing conditions and risk profile for the event

---

#### 8.3.4 TIME PERIOD AND FREQUENCY FOR EVENT ASSESSMENT AND FOLLOW-UP

The occurrence of an AE or SAE may come to the attention of study personnel from review of electronic medical record data or from a study participant presenting for medical care within the embedded PCP or the oncology model.

All AEs, not otherwise precluded per the protocol, will be captured on the appropriate CRF. Information to be collected includes event description, time of onset, clinician’s assessment of severity, relationship to study procedures (assessed only by those with the training and authority to make a diagnosis), and



time of resolution/stabilization of the event. All AEs occurring while on study will be documented appropriately regardless of relationship. All AEs will be followed to adequate resolution.

Any medical or psychiatric condition that is present at the time that the participant is screened will be considered as baseline and not reported as an AE. However, if the study participant's condition deteriorates at any time during the study, it will be recorded as an AE.

Changes in the severity of an AE will be documented to allow an assessment of the duration of the event at each level of severity to be performed. Documentation of onset and duration of each episode will be maintained for AEs characterized as intermittent.

The Principal Investigator and/or project manager will record events with start dates occurring any time after informed consent is obtained until 7 (for non-serious AEs) or 30 days (for SAEs) after the last day of study participation.

---

#### 8.3.5 ADVERSE EVENT REPORTING

The Principal Investigator will report AEs that are determined to be unanticipated problems to the KPSC IRB within 10 business days of discovery and at the continuing review period.

---

#### 8.3.6 SERIOUS ADVERSE EVENT REPORTING

The Principal Investigator will be responsible for conducting an evaluation of a SAE based on the unanticipated problem analysis and shall report the results of such evaluation to the NIH and the KPSC IRB as soon as possible, but no event later than 10 working days after the investigator first learns of the event. In the event of the death of a study participant, reporting will fall under one of the following pathways:

1. Death of an intervention study participant + unanticipated problem determination – the Principal Investigator will report to the Institutional Review Board within 1 business day of discovery and at continuing review
2. Death of an intervention study participant + NOT an unanticipated problem – the Principal Investigator will report at continuing review
3. Death of a non-interventional study participant – the event is not reportable

---

#### 8.3.7 REPORTING EVENTS TO PARTICIPANTS

N/A

---

### 8.3.8 EVENTS OF SPECIAL INTEREST

N/A

---

### 8.3.9 REPORTING OF PREGNANCY

N/A

---

## 8.4 UNANTICIPATED PROBLEMS

---

### 8.4.1 DEFINITION OF UNANTICIPATED PROBLEMS

As per the definition provided by the KPSC IRB we will consider any AE under this protocol a reportable UP if the event meets all three criteria:

- Unexpected in nature, severity, or frequency
- Related or possibly related to participation in the research
- Suggests greater risk to participants or others than previously known

Corrective actions or changes that may be considered in response to a UP are:

- Modification of inclusion or exclusion criteria to mitigate the newly identified risks
- Implementation of additional safety monitoring procedures
- Suspension of consenting/enrollment of new participants or halting of study procedures for consented/enrolled participants
- Modification of informed consent documents to include a description of newly recognized risks
- Provision of additional information about newly recognized risks to previously consented/enrolled participants

---

### 8.4.2 UNANTICIPATED PROBLEMS REPORTING

The Principal Investigator will report UPs to the reviewing KPSC IRB. The UP report will include the following information:

- Protocol identifying information: protocol title and number, Principal Investigator's name, and the IRB project number
- A detailed description of the event, incident, experience, or outcome
- An explanation of the basis for determining that the event, incident, experience, or outcome represents an UP
- A description of any changes to the protocol or other corrective actions that have been taken or are proposed in response to the UP

To satisfy the requirement for prompt reporting, UPs will be reported using the following timeline:

- UPs that are SAEs will be reported to the IRB and the study funding agency within 10 business days of the investigator becoming aware of the event; within one day in the event of the death of a participant if determined a UP.
- Any other UP will be reported to the IRB and to the funding agency within 10 business days of the investigator becoming aware of the problem

---

#### 8.4.3 REPORTING UNANTICIPATED PROBLEMS TO PARTICIPANTS

N/A

## 9 STATISTICAL CONSIDERATIONS

### 9.1 STATISTICAL HYPOTHESES

#### **Primary Endpoint(s):**

To determine the efficacy of the ePCP model relative to the usual care group on receipt of recommended care. We will construct a composite measure of guideline-recommended cancer surveillance and age-appropriate preventive care services, identifying utilization from EMR data. We hypothesize that patients in the ePCP model will have superior receipt of recommended care compared to usual care.

#### **Secondary Endpoint(s):**


- 1) To compare patient-reported outcomes (PROs) including patient-provider communication, coordination of survivorship care, and quality of life between the two models. We hypothesize that patients in the ePCP model will perceive significantly better care communication, self-efficacy, and confidence in delivery of survivorship care compared to usual care, as collected by PRO surveys.
- 2) To compare utilization of unplanned hospitalizations, use of urgent care, and receipt of non-recommended cancer surveillance services between the two models. We hypothesize that use of unplanned and non-recommended care will be significantly less in the ePCP model compared to usual care.

### 9.2 SAMPLE SIZE DETERMINATION


We computed minimum detectable effect sizes for fixed power (0.8) and level of significance (0.025), using superiority hypotheses for the primary (receipt of recommended care) and secondary PROs. As we expect the ePCP model to provide more comprehensive care, superiority hypotheses for these outcomes are warranted.

**Receipt of recommended care (Aim 1):** Expected prevalence of surveillance items in the first year of breast cancer survivorship range from 15% for assessment of bone health to 80% for mammography.<sup>118</sup> For CRC surveillance, a meta-analysis has shown that adherence ranged from 18-61% for colonoscopy and 17-71% for CEA tests. For preventive measures, Snyder et al. found that 30% of breast cancer survivors received recommended CRC screening, 32% received lipid testing, and 53% received an influenza vaccine. We expect to have an enrolled sample of approximately 3750 survivors with 268 patients at each site (5 embedded PCP sites, 9 usual care). Using PASS 14, we modeled a range of expected values from 15% to 80% and found that a study this size would provide 90% power to reject the null hypothesis when the mean difference between groups is between 4.2% and 5.5% with no clustering within centers (ICC=0) (Figure 1).

Based on similar KPSC studies, we expect observations from the same medical center to be only minimally correlated. We have modeled additional power analyses to account for this using an intraclass correlation (ICC) of 0.01. Using the same approach as above, we found that a study this size would provide 90% power to reject the null hypothesis when the mean difference between groups is between 7.5% and 10.5% with an ICC of 0.01 (Figure 2). Thus, we are well-powered to answer our primary research question.




**Figure 1.** Primary outcome (receipt of recommended care) power. A=alpha, the probability of rejecting a true null hypothesis ( $H_0$ :  $\text{Prevalence}_{\text{PCP}} \leq \text{Prevalence}_{\text{usualcare}}$ ); Pwr= power; K1= number of clusters in the PCP group (K2 in usual care group); M1=number of survivors per cluster in PCP group (M2 in usual care group); D0: margin of superiority; ICC= intraclass correlation; T= test (Z-test unpoiled)



**Figure 2.** Primary outcome with ICC=0.01

**PROs (Aim 2):** For our survey, we anticipate a 30% response rate with an estimated 1125 responders. For our secondary outcome of patient-provider communication, from prior studies using the MEPS measure we expect that 32% of patients will report high-quality communication, with responses to individual questions ranging from 29% to 62%. Using PASS 14, we calculated that our sample size of 1125, assuming 500 intervention and 625 control, would provide 80% power to reject the null hypothesis when the mean difference between groups exceeds 9.5% assuming there is no clustering within medical center (Figure 3).

For a minimal ICC of 0.01, using PASS 14, we calculated that this sample size would provide 80% power to reject the null hypothesis when the mean difference between groups exceeds 13.4% (Figure 4). Thus, we are well-powered to detect differences between the groups for our secondary outcome.



**Figure 3.** Secondary outcome (patient reported patient-provider communication) power.

### 9.3 POPULATIONS FOR ANALYSES

The analyses will follow an intent-to-treat (ITT) strategy, i.e. analyses will include all survivors in the groups to which their medical centers were assigned, regardless of their individual adherence with the model.

### 9.4 STATISTICAL ANALYSES

#### 9.4.1 GENERAL APPROACH

Descriptive statistics will be calculated prior to conducting the primary analyses, and these will be compared between the pre- and post-implementation periods to assess for potential imbalances in patient characteristics. We will check for data consistency and assumptions (e.g., normality of responses). Any data transformation or alternative methods necessary to analyze the data will be determined by examining the data structure. Individual survivors will be the unit of analysis. The analyses will follow an ITT strategy, i.e. analyses will include all survivors in the groups to which their medical centers were randomly assigned, regardless of their individual adherence with the model. We will include examination

of differences by gender and race/ethnicity for all analyses. We anticipate minimal missing data, based on our prior work using the EMR; if we encounter missing data, we will use the missing indicator method as this approach does not produce biased estimates for trials and is suitable for ITT analyses.<sup>119</sup> As a sensitivity analysis, we will compare results to complete case analysis and, for primary outcomes, multiple imputation for the missing covariate data. We expect the latter to closely match the results from the missing indicator method.

#### 9.4.2 ANALYSIS OF THE PRIMARY ENDPOINT(S)

For utilization data in both the pre- and post-periods, we will construct a composite measure of recommended care for cancer surveillance and for preventive care, based on recommendations from the ACS survivorship guideline. Cancer surveillance for early-stage breast cancer includes annual mammography and 2-4 physician visits with history and physical (H&P) for the first 3 years; CRC surveillance includes colonoscopy within first year, annual chest/abdominal/pelvic CT, and 2-4 H&P visits and CEA lab tests annually for the first 3 years. Preventive care measures will include appropriate cancer screening (other than for surveillance of primary cancer) for CRC, cervical, breast, and lung cancers, vaccinations (flu, pneumococcal), counseling for smoking cessation, and management of hyperlipidemia and hypercholesterolemia. These recommendations will be tailored to individual age, cancer type, health behaviors, and prior care (e.g., for breast cancer survivors  $\geq 50$ , colonoscopy every 10 years or annual FIT Kit for CRC cancer screening). We will use validated algorithms from our prior work based on CPT and ICD codes to identify use of services.<sup>17,31,32,99-102</sup> We will identify receipt of recommended care for up to 36 months, starting 6-12 months after cessation of active treatment (see Study Timeline, section 2.7 of the Study Record). Recommended care will be assessed in 12-month periods, with a binary outcome of received/did not receive for recommended cancer surveillance and for preventive care. We will also compare rates of individual services (e.g., rates of vaccination). Survivors with a pattern indicative of recurrence, based on validated recurrence algorithms,<sup>103,104</sup> will be excluded from analysis of recommended cancer care. We will identify use of services from oncology, embedded PCP, and the member's assigned KPSC PCP to evaluate the impact of the model on utilization of services.

For the primary outcomes, we will use the DID framework (see equation 1) with different link functions to address the different types of outcomes (continuous, categorical, or counts). Outcome measurements are denoted  $y_{ijkt}$ , where  $i$  indexes the individual patients,  $j$  indexes the providers with whom patients had follow-up visit at site  $k$ , and during study time period  $t$ . The first two parameters on the right-hand side of equation 1 denote a random clinician intercept ( $\delta_{0j}$ ) to represent the baseline variation between physicians in receiving recommended care, and  $\delta_1$  denotes the level difference for the Oncology-led model of care at the beginning of the post-implementation follow-up period. The other two DID parameters  $\delta_2$  and  $\delta_3$  specify the average difference between Embedded-PCP and Oncology-led models at baseline and the difference-in-differences, the latter of which is the main quantity of interest. The remaining parameters correspond to fixed effects for respective patient, provider and site characteristics.

**Equation 1:**  $f(E(y_{ijkt})) = \delta_{0j} + \delta_1 I(\text{post}=1) + \delta_2 I(\text{Embedded PCP}=1) + \delta_3 I(\text{post}=1) I(\text{Embedded PCP}=1) + \alpha_p + \alpha_t + X_i \alpha + X_j \beta + X_k \gamma$

#### 9.4.3 ANALYSIS OF THE SECONDARY ENDPOINT(S)

The main outcomes for the Aim 2 survey will be analyzed using the general modeling approach outlined in Equations 2 and 3. In this case,  $y_{ijk}$  could denote the probability that patient  $i$  reports ‘high quality’ patient-provider communication during study period  $k$ . Unlike the primary outcome in Aim 1, we do not intend to collect repeated measures of these patient-reported outcomes, and so may only include provider-level random effects. Other parameters in the model may be interpreted in the same manner as described above. We will similarly examine possible time-by-treatment interactions to assess whether, for example, differences in patient-provider communication change over time.

**Equation 2.**  $f(E(y_{ijkt})) = \delta_{0j} + \delta_1 I(\text{Embedded PCPI}=1) + X_{ijk} \beta + X_{it} \gamma$   
**Equation 3.**  $\delta_{0j} \sim \text{Normal}(0, \sigma_0^2)$

For Aim 3, we will use CPT and ICD codes to identify hospitalizations related to cancer, using primary ICD code to identify reason for admission, as well as use of urgent care and use of non-recommended care. Non-recommended care measures are identified from relevant guidelines (e.g., ASCO Choosing Wisely, ACS Survivorship).<sup>115-117</sup> For early-stage breast cancer, this includes serum tumor markers and high-intensity imaging (e.g., CT and PET scans); for CRC, PET scans.

#### 9.4.4 SAFETY ANALYSES

N/A

#### 9.4.5 BASELINE DESCRIPTIVE STATISTICS

Demographic information (e.g., age, race, ethnicity, census-level education and income) and clinical information (e.g., date of cancer diagnosis, age at diagnosis, disease stage, cancer treatment, endocrine therapy) will be collected from the EMR for all participants and compared between intervention and control groups.

#### 9.4.6 PLANNED INTERIM ANALYSES

N/A

#### 9.4.7 SUB-GROUP ANALYSES

We will test for differences by gender and race/ethnicity between the intervention and control groups on utilization and patient-reported outcomes. Prior studies of utilization in cancer survivors do not suggest or negate significant differences in intervention effect among subgroups; thus, it is important for our study to examine potential differences.

---

#### 9.4.8 TABULATION OF INDIVIDUAL PARTICIPANT DATA

We will not tabulate individual participant data.

---

#### 9.4.9 EXPLORATORY ANALYSES

We will explore PCP knowledge of and confidence in providing survivorship care, comparing the embedded PCPs to PCPs practicing at centers non-randomly assigned to the usual care model. We will compare embedded PCPs to usual care PCPs using 2-sided chi-square tests.

### 10 SUPPORTING DOCUMENTATION AND OPERATIONAL CONSIDERATIONS

#### 10.1 REGULATORY, ETHICAL, AND STUDY OVERSIGHT CONSIDERATIONS

---

##### 10.1.1 INFORMED CONSENT PROCESS

---

###### 10.1.1.1 CONSENT/ASSENT AND OTHER INFORMATIONAL DOCUMENTS PROVIDED TO PARTICIPANTS

For this study, a waiver for informed consent will be required to ensure a scientifically valid comparison of the ePCP model to current usual care. We do not foresee any risks to potential participants by not directly consenting them to participate in the trial. This study poses minimal risk to the patient because survivorship cancer care will be performed as part of routine medical care, and both the ePCP and the oncology-based model for survivorship cancer care are consistent with guidelines from professional societies and organizations with regard to the delivery of safe, appropriate and usual care practices. We also believe that such a waiver of consent is not only widely recognized as necessary and appropriate, but also scientifically necessary and ethically justifiable under the “Common Rule” for protection of research participants (45 CFR 46.116d).

Patients will be recruited for PRO surveys under Aim 2. We will obtain written or oral consent from all participants for participation in the surveys. Survey consents and opt out options will be built into an electronic template as part of the initial survey screens. Consent forms describing in detail the survey procedures, and risks will be given to the participant and written or oral documentation of informed consent will be completed prior to completing the survey. The following consent materials are submitted with this protocol [none at this time; however, all written or oral consenting procedures will be submitted to our IRB prior to conducting any patient contact for the surveys under this Aim.



For utilization data captured under Aim 3 from the EMR, individual patients will not be recruited or consented (falling under the waiver of informed consent for Aim 1).

---

#### 10.1.1.2 CONSENT PROCEDURES AND DOCUMENTATION

For Aims 1 and 3, consenting procedures are not applicable.

For Aim 2, consent forms describing in detail the survey procedures, and risks will be given to the participant, and written or oral documentation of informed consent will be completed prior to completing the survey.

---

#### 10.1.2 STUDY DISCONTINUATION AND CLOSURE

This study may be temporarily suspended or prematurely terminated if there is sufficient reasonable cause. Written notification, documenting the reason for study suspension or termination, will be provided by the suspending or terminating party to study participants, investigator, funding agency, and regulatory authorities. If the study is prematurely terminated or suspended, the Principal Investigator (PI) will promptly inform study participants, the KPSC IRB, and sponsor/funding agency and will provide the reason(s) for the termination or suspension. Study participants will be contacted, as applicable, and be informed of changes to study visit schedule.

Circumstances that may warrant termination or suspension include, but are not limited to:

- Determination of unexpected, significant, or unacceptable risk to participants
- Demonstration of efficacy that would warrant stopping
- Insufficient compliance to protocol requirements
- Data that are not sufficiently complete and/or evaluable
- Determination that the primary endpoint has been met
- Determination of futility

The study may resume once concerns about safety, protocol compliance, and data quality are addressed, and satisfy the funding agency, KPSC IRB, or other relevant regulatory or oversight bodies (OHRP, DSMB).]

---

#### 10.1.3 CONFIDENTIALITY AND PRIVACY

Participant confidentiality will be maintained at all times. Each survey participant will be given a unique alpha-numeric identification number. No data collection form will be linked to an actual participant name. Any document linking the participant name to an identification number will be kept in a locked file separate from data collection files. All records from the study will be kept in locked files within the KPSC Department of Research and Evaluation. Identifying information will be removed from the final data sets

and stored in a linked file to which only the Principal Investigator or designee has access. Individual participants will not be identified in any reports. Additionally, study reports will be aggregated so that individual participants are not identified. All investigators and research staff at participating sites will be required to maintain up-to-date training in human subject protection and good clinical practice through Collaborative Institutional Training Initiative (CITI; <https://www.citiprogram.org/default.asp>) and HIPAA training. Additional security precautions include encryption, digital certification, audit logs, and firewall protection.

The study monitor, other authorized representatives of the sponsor or funding agency, or representatives of the KPSC IRB may inspect all documents and records required to be maintained by the investigator, including but not limited to, medical records (office, clinic, or hospital) for the participants in this study. The clinical study site will permit access to such records. At the end of the study, all records will continue to be kept in a secure location for as long a period as dictated by the KPSC IRB.

**Measures Taken to Ensure Confidentiality of Data Shared per the NIH Data Sharing Policies.** The Principal Investigator will ensure all mechanisms used to share data will include proper plans and safeguards for the protection of privacy, confidentiality, and security for data dissemination and reuse (e.g., all data will be thoroughly de-identified and will not be traceable to a specific study participant; see Section 10.1.4 for details). Plans for archiving and long-term preservation of the data will be implemented, as appropriate. A Certificate of Confidentiality is not required for this study.

---

#### 10.1.4 FUTURE USE OF STORED SPECIMENS AND DATA

We are committed to the open and timely dissemination of study data and research outcomes to the broader community of health care providers, investigators, patients, health care systems, and public health personnel. The investigators agree to abide by the principles for sharing research resources as described by the National Cancer Institute's Cancer Moonshot public access and data sharing policy. Publications based on the study will adhere to the NIH Public Access Policy. Study data will be de-identified and made available for use in subsequent studies by other investigators to the extent feasible, pursuant to details as outlined below. We will invite other investigators to submit ancillary study proposals using the data from the study.

The study will also produce deliverables that will be freely available to the cancer care community including protocols for the interventions as well as EMR algorithms (e.g., risk-stratification), and clinician and patient educational tools. We will engage in presentation and publication of findings, and have structured each Aim to have publishable results, with an expectation of multiple opportunities for scientific presentations and manuscripts.

Our plan to support efforts by other researchers to replicate this study in other oncology settings and practices, as well as for other healthcare systems that wish to test and/or implement the resulting products, includes: 1) sharing of a complete, detailed study protocol; 2) sharing of intervention tools developed during the study; and 3) sharing of datasets. Each of these activities will ensure transparency

as well as replication so that other researchers will be able to apply the same study procedures, measures and analytic approaches to similar or novel populations. Our sharing of de-identified data will allow researchers to consider additional or associated questions or as a basis for collecting new data using this study design.

---

#### 10.1.5 KEY ROLES AND STUDY GOVERNANCE

| <b>Principal Investigator</b>            |
|--|
| Erin E. Hahn, PhD, MPH, Res. Scientist I |
| Kaiser Permanente Southern California    |
| Department of Research & Evaluation      |
| 100 S. Los Robles, 2 <sup>nd</sup> Floor |
| 626-564-3505                             |
| Erin.E.Hahn@kp.org                       |

We have assembled an Advisory Board with expertise in complementary areas to the study team. We will also include patient members who have experienced cancer, recruited through the KPSC Regional Member Advisory Committee. The role of the Board will be to provide input and guidance on the overall study design and progress, including refinement of model components (e.g., PCP training, health IT alerts), measurement of outcomes, and discussion of results. We will convene quarterly advisory board meetings each year. We will rely on stakeholder engagement strategies to ensure that our work is relevant and responsive, following the general approach of the Six Stage model for engagement.

---

#### 10.1.6 SAFETY OVERSIGHT

The Data and Safety Monitoring Plan (DSMP) for this study includes monitoring recruitment procedures and potential adverse events resulting from data collection (observational data from clinical and administrative databases, and survey data) and from the intervention. There are minimal physical, psychological, or other risks associated with the proposed research based on the Common Rule, 45 CFR 46.116d, because this study involves the use of routine care practices, no investigational drugs or devices, and poses little risk of physical or psychological harm to patients in either the embedded PCP or oncology model (standard of care) arm under Aim 1.

We have appointed a Data and Safety Monitoring Board (DSMB) consisting of a biostatistician, a cancer epidemiologist, and an oncologist. The DSMB will meet with the investigators during the start-up phase to approve the final study protocol. Yearly meetings will occur via conference calls to monitor recruitment and adverse events.

---

#### 10.1.7 CLINICAL MONITORING

N/A

---

#### 10.1.8 QUALITY ASSURANCE AND QUALITY CONTROL

Quality control (QC) procedures will be implemented as follows:

**Informed consent** ---

- Aim 1, not applicable. The medical center sites will not consent participants under Aim 1.
- Aim 2, consenting for the PRO survey will be managed by the research study team (study staff will review both the documentation of the consenting process as well as a percentage of the completed consent documents). This review will evaluate compliance with GCP, accuracy, and completeness. Feedback will be provided to the study team to ensure proper consenting procedures are followed.
- Aim 3, not applicable.

**Source documents and the electronic data** --- Data for this study are obtained from clinical and administrative databases (Aim 1) and participant self-report for the surveys (Aim 2). For utilization data under Aim 3, we will use a similar approach to Aim 1 – e.g., CPT and ICD codes will be used to identify hospitalizations related to cancer, use of urgent care, and use of non-recommended care. Non-recommended care measures will be identified from relevant guidelines (e.g., ASCO Choosing Wisely, ACS Survivorship). We will collect data in order to initiate participation identification at all sites in Years 1 and 2 (Aim 1); we will collect PRO survey data in Years 2-4 (Aim 2); we will collect utilization data in Year 5 (Aim 3).

**Intervention Fidelity** — We will assess compliance with the study protocol monthly, with review of relevant data. We will collect utilization data for services with the ePCP, the treating oncology team, and with the patient's assigned KPSC PCP. We will check for patterns of utilization indicative non-compliance with the ePCP protocol (e.g., no or limited visits with the ePCP, continued use of oncology for patients in the ePCP arm). We will have regular interactions with the study sites to discuss and review these data.

Clinical EMR data on utilization has a high degree of accuracy but will be reviewed at regular intervals to ensure quality and accuracy (e.g., chart review of outliers). Interim data checks will be done in aggregate, with means, median, and range checks performed quarterly.

**Protocol Deviations** – Throughout the study phases, we will work closely with our Institutional Review Board, clinical chiefs, and operational leaders to ensure patient safety, with regular review and reporting of adverse events and protocol deviations.

Should independent monitoring become necessary, the Principal Investigator will provide direct access to all trial related sites, source data/documents, and reports for the purpose of monitoring and auditing by the sponsor/funding agency, and inspection by local and regulatory authorities.

---

### 10.1.9 DATA HANDLING AND RECORD KEEPING

---

#### 10.1.9.1 DATA COLLECTION AND MANAGEMENT RESPONSIBILITIES

Demographic information (e.g., age, race, ethnicity, census-level education and income) and clinical information (e.g., date of cancer diagnosis, age at diagnosis, disease stage, cancer treatment, endocrine therapy) will be collected from the EMR for all participants (Aim 1) by study programmers under the oversight of Dr. Hahn. Self-reported demographics will be collected from those who participate in the PRO survey (Aim 2). We will collect clinician variables such as clinician gender, race, and years in practice from the EMR.

Clinical EMR data on utilization (Aim 3) will be reviewed at regular intervals to ensure quality and accuracy (e.g., chart review of outliers). Interim data checks will be done in aggregate, with means, median, and range checks performed quarterly.

All investigators and research staff at participating sites will be required to maintain up-to-date training in human subject protection and good clinical practice through Collaborative Institutional Training Initiative (CITI; <https://www.citiprogram.org/default.asp>) and Health Insurance Portability and Accountability Act (HIPAA) training. Additional security precautions include encryption, digital certification, audit logs, and firewall protection.

---

#### 10.1.9.2 STUDY RECORDS RETENTION

Study documents will be retained for a minimum of 2 years after cessation of study analyses. These documents should be retained for a longer period, however, if required by local regulations or additional studies using these data and resources.

---

### 10.1.10 PROTOCOL DEVIATIONS

This protocol uses the KPSC definition of protocol deviation: a protocol deviation is a departure from the IRB approved research plan that – a) does not place the safety, rights, or welfare of one or more study participants at risk AND b) does not impact the integrity of the study. As a result of deviations, corrective actions will be developed by the site and implemented promptly.

It will be the responsibility of the Principal Investigator to use continuous vigilance to identify and report deviations at the next continuing review period of identification of the protocol deviation. All deviations will be addressed in study source documents, reported to the KPSC IRB for review per their policies. The site investigator will be responsible for knowing and adhering to the KPSC IRB requirements.

---

### 10.1.11 PUBLICATION AND DATA SHARING POLICY

This study will be conducted in accordance with the following publication and data sharing policies and regulations:

NIH Public Access Policy, which ensures that the public has access to the published results of NIH funded research. It requires scientists to submit final peer-reviewed journal manuscripts that arise from NIH funds to the digital archive PubMed Central upon acceptance for publication.

This study will comply with the NIH Data Sharing Policy and Policy on the Dissemination of NIH-Funded Clinical Trial Information and the Clinical Trials Registration and Results Information Submission rule. As such, this trial will be registered at ClinicalTrials.gov, and results information from this trial will be submitted to ClinicalTrials.gov. In addition, every attempt will be made to publish results in peer-reviewed journals. Data from this study may be requested from other researchers up to 5 years after the completion of the primary endpoint by contacting Dr. Erin E. Hahn. Considerations for ensuring confidentiality of these shared data are described in **Section 10.1.3**.

**Details of data sharing include but are not limited to:**

### **1. Sharing of a complete, detailed study protocol and publications**

We will ensure our study protocol is registered on ClinicalTrials.gov prior to initiation of enrollment procedures. At the end of the study, we will provide an updated, detailed study protocol describing all aspects of the study including the following: a) description of the study population, risk-stratification, sampling methods and resulting study sample; b) study design; recruitment and enrollment procedures; and primary care intervention components and implementation; c) data collection including programming algorithms to extract CPT-based utilization data; and d) copies of survey measures and data collection schedules and procedures. Our grant application becomes the initial draft of this protocol. This protocol will be updated at the end of the study planning phase to ensure that it provides a concrete, detailed, and specific step-by-step manual to allow replication. For reproducibility and for full transparency, any code or tools developed will be published along with the manuscript either through the journal website and/or GitHub or Open Science Framework website, which is hosted by the Center for Open Science (funded in part by the NIH and the National Science foundation) and is open to the public. We will also seek to publish this protocol in a relevant scientific journal. This and other publications generated from this study will:

1. Be deposited in PubMed Central with proper tagging of metadata after acceptance by a journal;
2. Will be published under the Creative Commons Attribution 4.0 Generic License or an equivalent license, or otherwise dedicated to the public domain;
3. To the extent feasible, underlying primary data will be shared simultaneously with the publication.

### **2. Sharing of de-identified data for replication and additional studies**

For this study, the Investigators will provide these research data in a controlled manner to outside researchers who are willing to enter into formal research relationships to ensure that the data will be used for scientific purposes in the public interest, that patient privacy will be protected, and that all other risks

to participants will be minimized. All such requests will be reviewed by the Investigators for scientific merit, human subjects considerations, and KPSC legal obligations. Primary data will reside locally within the Kaiser Permanente Department of Research and Evaluation. After approval of the request, a data sharing agreement will be created, approved, and signed. Conditions may be placed on the use of the data, including but not limited to, no distribution to third parties, a KPSC researcher included on the study team proper acknowledgement and citation of the data providers (as indicated in the data sharing agreement) and the NIH grant funding, exclusive use by the data recipient in connection with a specific research project, for which the recipient has sole responsibility and which is explicitly described, and agreement not to use the data in any effort to establish the identity of the study subjects. The data recipient will be subject to applicable federal, state, and local laws or regulations and institutional policies providing additional protections for human subjects.

De-identification of data: In order to protect the privacy rights of participants and confidentiality of their data, we will de-identify all data according to the standards set forth in the Health and Human Services Regulations for the Protection of Human Subjects; primary data will also be stripped of identifiers according to the HIPAA Privacy Rule. With input from the KPSC IRB, we will assess consent materials to determine whether data may be shared as contemplated in this policy.

If de-identified data can be shared, the following process and procedures is to be followed:

1. Requests will be evaluated individually by the Principal Investigator of this study in conjunction with the Senior Director of Research and/or the Executive Committee of the Department of Research and Evaluation.
  - a. Requests must be from a qualified, doctorate-level researcher;
  - b. Requests must have a stated, achievable purpose and detailed plan that is of value to the clinical and/or research community.
2. A data use agreement and/or data transfer agreement must be executed for all requests. This must include:
  - a. What data elements are to be included;
  - b. How the data will be used;
  - c. Details of any subsequent disclosure and prohibition of additional disclosure without an amendment;
  - d. How long the data can be used;
  - e. What will happen to the data when the project is complete;
  - f. Opportunity for review and comment.
3. De-identified datasets to be made available under this Plan must:
  - a. Follow the definitions of de-identification as outlined by Health and Human Services Regulations for the Protection of Human Subjects and the HIPAA Privacy Rule;
  - b. Use relative dates and round birth dates/ages;
  - c. De-identify entity/site of care to be to the extent possible;
  - d. Only include data to support publication.
4. An approval process will be in place, with required approvals from:
  - a. SCPMG practice leader (typically Regional Chief of relevant specialty)

- b. SCPMG Area Medical Director
  - c. Senior Director of Research, Department of Research and Evaluation
5. There will be no associated fees.

### 10.1.12 CONFLICT OF INTEREST POLICY

Any actual conflict of interest of persons who have a role in the design, conduct, analysis, publication, or any aspect of this trial will be disclosed and managed. Furthermore, persons who have a perceived conflict of interest will be required to have such conflicts managed in a way that is appropriate to their participation in the design and conduct of this trial. The study leadership in conjunction with the National Cancer Institute (NCI) has established policies and procedures for all study group members to disclose all conflicts of interest and will establish a mechanism for the management of all reported dualities of interest.

### 10.2 ADDITIONAL CONSIDERATIONS

N/A

### 10.3 ABBREVIATIONS AND SPECIAL TERMS

|        |  |
|--------|--|
| AE     | Adverse Event  |
| ACS    | American Cancer Institute                                      |
| AHRQ   | Agency for Healthcare Research and Quality                     |
| ASCO   | American Society for Clinical Oncology                         |
| CAHPS  | Consumer Assessment of Healthcare Provider and Systems         |
| CEA    | Carcinoembryonic Antigen                                       |
| CFR    | Code of Federal Regulations                                    |
| CITI   | Collaborative Institutional Training Initiative                |
| CMP    | Clinical Monitoring Plan                                       |
| COC    | Certificate of Confidentiality                                 |
| CRC    | Colorectal Cancer  |
| CRF    | Case Report Form   |
| CVD    | Cardiovascular Disease   |
| DHHS   | Department of Health and Human Services                        |
| DSMB   | Data Safety Monitoring Board                                   |
| EORTC  | European Organisation for the Research and Treatment of Cancer |
| EMR    | Electronic Medical Record(s)                                   |
| ePCP   | Embedded Primary Care Physician                                |
| FITKit | Fecal Immunochemical Test Kit                                  |
| GCP    | Good Clinical Practice   |
| H&P    | History and Physical   |
| HIPAA  | Health Insurance Portability and Accountability Act            |
| ICC    | Intracluster Correlation                                       |
| ICD    | International Classification of Diseases                       |



|        |   |
|--------|---|
| ICH    | International Conference on Harmonisation                 |
| IRB    | Institutional Review Board                                |
| IT     | Information Technology                                    |
| ITT    | Intent to Treat   |
| KPSC   | Kaiser Permanente Southern California                     |
| MEPS   | Medical Expenditure Panel Survey                          |
| MOP    | Manual of Procedures                                      |
| NCI    | National Cancer Institute                                 |
| NIH    | National Institutes of Health                             |
| OHRP   | Office for Human Research Protections                     |
| PCORI  | Patient-Centered Outcomes Research Institute              |
| PET    | Positron Emission Tomography                              |
| PHI    | Protected Health Information                              |
| PO     | Project Officer   |
| PROs   | Patient-Reported Outcomes                                 |
| PROMIS | Patient-Reported Outcomes Measurement Information Systems |
| QA     | Quality Assurance   |
| QC     | Quality Control   |
| RCT    | Randomized-Controlled Trial                               |
| SAE    | Serious Adverse Event                                     |
| SAP    | Statistical Analysis Plan                                 |
| SAS    | Statistical Analysis Software                             |
| SCPs   | Survivorship Care Plans                                   |
| SCPMG  | Southern California Permanente Medical Group              |
| SMC    | Safety Monitoring Committee                               |
| SO     | Safety Officer  |
| UK     | United Kingdom  |
| UP     | Unanticipated Problem                                     |
| URL    | Uniform Resource Locator                                  |
| US     | United States   |

## 10.4 PROTOCOL AMENDMENT HISTORY

[illegible]

## 11 REFERENCES

1. Bluethmann SM, Mariotto AB, Rowland JH: Anticipating the "Silver Tsunami": Prevalence Trajectories and Comorbidity Burden among Older Cancer Survivors in the United States. *Cancer Epidemiol Biomarkers Prev* 25:1029-36, 2016
2. Parry C, Kent EE, Mariotto AB, et al: Cancer Survivors: A Booming Population. *Cancer Epidemiology Biomarkers & Prevention* 20:1996-2005, 2011
3. Denlinger CS, Carlson RW, Are M, et al: Survivorship: introduction and definition. Clinical practice guidelines in oncology. *Journal of the National Comprehensive Cancer Network : JNCCN* 12:34-45, 2014
4. Ganz PA: Monitoring the physical health of cancer survivors: a survivorship-focused medical history. *J Clin Oncol* 24:5105-11, 2006
5. Ganz PA: Late effects of cancer and its treatment. *Semin Oncol Nurs* 17:241-8, 2001
6. Kline RM, Arora NK, Bradley CJ, et al: Long-Term Survivorship Care After Cancer Treatment - Summary of a 2017 National Cancer Policy Forum Workshop. *J Natl Cancer Inst* 110:1300-1310, 2018
7. Hewitt M, Greenfield S, Stovall E: From Cancer patient to cancer survivor: Lost in transition, Institute of Medicine and National Research Council 2005
8. Adler N, Page A: Cancer care for the whole patient: Meeting psychosocial health needs - National Institutes of Medicine Committee on Psychosocial Services to Cancer Patients/families in a Community Setting. Washington D.C, National Academies Press, 2008
9. Rehis R BE, Arvey SR, et al. The Essential Elements of Survivorship Care: A Livestrong Brief. The National Cancer Survivorship Resource Center, American Cancer Society, and The GW Cancer Institute: [https://www.livestrong.org/sites/default/files/what-we-do/reports/Essential-Elements-Definitions\\_Recommendations.pdf](https://www.livestrong.org/sites/default/files/what-we-do/reports/Essential-Elements-Definitions_Recommendations.pdf).
10. Jacobs LA, Shulman LN: Follow-up care of cancer survivors: challenges and solutions. *Lancet Oncol* 18:e19-e29, 2017
11. Nekhlyudov L, Mollica MA, Jacobsen PB, et al: Developing a Quality of Cancer Survivorship Care Framework: Implications for Clinical Care, Research and Policy. *JNCI: Journal of the National Cancer Institute*, 2019
12. American Society of Clinical Oncology. Guidelines on Survivorship Care. Available: <https://www.asco.org/practice-guidelines/cancer-care-initiatives/prevention-survivorship/survivorship-compendium-0>.
13. National Comprehensive Cancer Network Guidelines. Available: [https://www.nccn.org/professionals/physician\\_gls/f\\_guidelines.asp](https://www.nccn.org/professionals/physician_gls/f_guidelines.asp).
14. Viswanathan M, Halpern M, T. SE, et al: Models of Cancer Survivorship Care: Agency for Healthcare Reserach and Quality. Rockville, MD, 2014
15. Howell D, Hack T, Oliver T, et al: Models of care for post-treatment follow-up of adult cancer survivors: a systematic review and quality appraisal of the evidence. *Journal of Cancer Survivorship* 6:359-71, 2012

16. Halpern MT, Viswanathan M, Evans TS, et al: Models of Cancer Survivorship Care: Overview and Summary of Current Evidence. *Journal of Oncology Practice* 11:e19-e27, 2014
  17. Hahn EE, Hays RD, Kahn KL, et al: Use of imaging and biomarker tests for posttreatment care of early-stage breast cancer survivors. *Cancer* 119:4316-24, 2013
  18. Keating NL, Landrum MB, Guadagnoli E, et al: Factors related to underuse of surveillance mammography among breast cancer survivors. *J Clin Oncol* 24:85-94, 2006
  19. Schapira MM, McAuliffe TL, Nattinger AB: Underutilization of mammography in older breast cancer survivors. *Med Care* 38:281-9, 2000
  20. Salz T, Weinberger M, Ayanian JZ, et al: Variation in use of surveillance colonoscopy among colorectal cancer survivors in the United States. *BMC Health Serv Res* 10:256, 2010
  21. Sisler JJ, Seo B, Katz A, et al: Concordance with ASCO guidelines for surveillance after colorectal cancer treatment: a population-based analysis. *J Oncol Pract* 8:e69-79, 2012
  22. Urquhart R, Folkes A, Porter G, et al: Population-based longitudinal study of follow-up care for patients with colorectal cancer in Nova Scotia. *Journal of oncology practice* 8:246-252, 2012
  23. Earle CC, Burstein HJ, Winer EP, et al: Quality of Non–Breast Cancer Health Maintenance Among Elderly Breast Cancer Survivors. *Journal of Clinical Oncology* 21:1447-1451, 2003
  24. Kunitake H, Zheng P, Yothers G, et al: Routine preventive care and cancer surveillance in long-term survivors of colorectal cancer: results from National Surgical Adjuvant Breast and Bowel Project Protocol LTS-01. *J Clin Oncol* 28:5274-9, 2010
  25. Earle CC, Neville BA: Under use of necessary care among cancer survivors. *Cancer* 101:1712-1719, 2004
  26. Weaver KE, Foraker RE, Alfano CM, et al: Cardiovascular risk factors among long-term survivors of breast, prostate, colorectal, and gynecologic cancers: a gap in survivorship care? *J Cancer Surviv* 7:253-61, 2013
  27. Johnson CB, Davis MK, Law A, et al: Shared Risk Factors for Cardiovascular Disease and Cancer: Implications for Preventive Health and Clinical Care in Oncology Patients. *Canadian Journal of Cardiology* 32:900-907, 2016
  28. Elston Lafata J, Simpkins J, Schultz L, et al: Routine surveillance care after cancer treatment with curative intent. *Med Care* 43:592-9, 2005
  29. Salloum RG, Hornbrook MC, Fishman PA, et al: Adherence to surveillance care guidelines after breast and colorectal cancer treatment with curative intent, 2012
  30. Panageas KS, Sima CS, Liberman L, et al: Use of high technology imaging for surveillance of early stage breast cancer. *Breast Cancer Res Treat* 131:663-70, 2012
  31. Hahn EE, Munoz-Plaza C, Wang J, et al: Anxiety, Culture, and Expectations: Oncologist-Perceived Factors Associated With Use of Nonrecommended Serum Tumor Marker Tests for Surveillance of Early-Stage Breast Cancer. *J Oncol Pract* 13:e77-e90, 2017
  32. Hahn EE, Tang T, Lee JS, et al: Use of posttreatment imaging and biomarkers in survivors of early-stage breast cancer: Inappropriate surveillance or necessary care? *Cancer* 122:908-16, 2016
  33. Cooper GS, Johnson CC, Lamerato L, et al: Use of guideline recommended follow-up care in cancer survivors: routine or diagnostic indications? *Med Care* 44:590-4, 2006
- NIH Behavioral and Social Intervention Clinical Trial Protocol Template v3.0 - 20180827

34. Cooper GS, Kou TD, Reynolds HL, Jr.: Receipt of guideline-recommended follow-up in older colorectal cancer survivors : a population-based analysis. *Cancer* 113:2029-37, 2008
35. Cooper GS, Yuan Z, Chak A, et al: Geographic and patient variation among Medicare beneficiaries in the use of follow-up testing after surgery for nonmetastatic colorectal carcinoma. *Cancer* 85:2124-31, 1999
36. Ramsey SD, Fedorenko C, Chauhan R, et al: Baseline Estimates of Adherence to American Society of Clinical Oncology/American Board of Internal Medicine Choosing Wisely Initiative Among Patients With Cancer Enrolled With a Large Regional Commercial Health Insurer. *J Oncol Pract* 11:338-43, 2015
37. Urquhart R, Folkes A, Babineau J, et al: Views of breast and colorectal cancer survivors on their routine follow-up care. *Current oncology (Toronto, Ont.)* 19:294-301, 2012
38. Shulman LN, Jacobs LA, Greenfield S, et al: Cancer care and cancer survivorship care in the United States: will we be able to care for these patients in the future? *J Oncol Pract* 5:119-23, 2009
39. Erikson C, Salsberg E, Forte G, et al: Future supply and demand for oncologists : challenges to assuring access to oncology services. *J Oncol Pract* 3:79-86, 2007
40. Yang W, Williams JH, Hogan PF, et al: Projected supply of and demand for oncologists and radiation oncologists through 2025: an aging, better-insured population will result in shortage. *J Oncol Pract* 10:39-45, 2014
41. Mayer DK, Alfano CM: Personalized Risk-Stratified Cancer Follow-Up Care: Its Potential for Healthier Survivors, Happier Clinicians, and Lower Costs. *JNCI: Journal of the National Cancer Institute* 111:442-448, 2019
42. Bilimoria KY, Ko CY, Tomlinson JS, et al: Wait times for cancer surgery in the United States: trends and predictors of delays. *Ann Surg* 253:779-85, 2011
43. Snyder CF, Frick KD, Kantsiper ME, et al: Prevention, screening, and surveillance care for breast cancer survivors compared with controls: changes from 1998 to 2002. *J Clin Oncol* 27:1054-61, 2009
44. Snyder CF, Earle CC, Herbert RJ, et al: Preventive care for colorectal cancer survivors: a 5-year longitudinal study. *J Clin Oncol* 26:1073-9, 2008
45. Snyder CF, Earle CC, Herbert RJ, et al: Trends in follow-up and preventive care for colorectal cancer survivors. *J Gen Intern Med* 23:254-9, 2008
46. Snyder CF, Frick KD, Herbert RJ, et al: Comorbid condition care quality in cancer survivors: role of primary care and specialty providers and care coordination. *J Cancer Surviv* 9:641-9, 2015
47. Wallner LP, Li Y, Furgal AKC, et al: Patient Preferences for Primary Care Provider Roles in Breast Cancer Survivorship Care. *J Clin Oncol* 35:2942-2948, 2017
48. Tarasenko YN, Miller EA, Chen C, et al: Physical activity levels and counseling by health care providers in cancer survivors. *Prev Med* 99:211-217, 2017
49. Haggstrom DA, Arora NK, Helft P, et al: Follow-up care delivery among colorectal cancer survivors most often seen by primary and subspecialty care physicians. *J Gen Intern Med* 24 Suppl 2:S472-9, 2009
50. Forsythe LP, Alfano CM, Leach CR, et al: Who provides psychosocial follow-up care for post-treatment cancer survivors? A survey of medical oncologists and primary care physicians. *J Clin Oncol* 30:2897-905, 2012

51. Demark-Wahnefried W, Pinto BM, Gritz ER: Promoting Health and Physical Function Among Cancer Survivors: Potential for Prevention and Questions That Remain. *Journal of Clinical Oncology* 24:5125-5131, 2006
52. Pearlstein KA, Basak R, Chen RC: Cardiovascular Care Among Cancer Survivors in the United States. *JNCI Cancer Spectrum* 2, 2018
53. Hudson SV, Ohman-Strickland PA, Bator A, et al: Breast and prostate cancer survivors' experiences of patient-centered cancer follow-up care from primary care physicians and oncologists. *J Cancer Surviv* 10:906-14, 2016
54. Luctkar-Flude M, Aiken A, McColl MA, et al: Are primary care providers implementing evidence-based care for breast cancer survivors? *Canadian family physician Medecin de famille canadien* 61:978-984, 2015
55. Radhakrishnan A, Li Y, Furgal AKC, et al: Provider Involvement in Care During Initial Cancer Treatment and Patient Preferences for Provider Roles After Initial Treatment. *Journal of Oncology Practice* 15:e328-e337, 2019
56. Del Giudice ME, Grunfeld E, Harvey BJ, et al: Primary care physicians' views of routine follow-up care of cancer survivors. *J Clin Oncol* 27:3338-45, 2009
57. Hahn EE, Munoz-Plaza CE, Schottinger JE, et al: Developing innovative models of care for cancer survivors: use of implementation science to guide evaluation of appropriateness and feasibility. *Support Care Cancer* 27:1737-1745, 2019
58. Hudson SV, Miller SM, Hemler J, et al: Adult Cancer Survivors Discuss Follow-up in Primary Care: 'Not What I Want, But Maybe What I Need'. *The Annals of Family Medicine* 10:418-427, 2012
59. Klabunde CN, Haggstrom D, Kahn KL, et al: Oncologists' perspectives on post-cancer treatment communication and care coordination with primary care physicians. *European Journal of Cancer Care* 26:e12628, 2017
60. Potosky AL, Han PK, Rowland J, et al: Differences between primary care physicians' and oncologists' knowledge, attitudes and practices regarding the care of cancer survivors. *J Gen Intern Med* 26:1403-10, 2011
61. Rubinstein EB, Miller WL, Hudson SV, et al: Cancer survivorship care in advanced primary care practices: A qualitative study of challenges and opportunities. *JAMA Internal Medicine*, 2017
62. Virgo KS, Lerro CC, Klabunde CN, et al: Barriers to breast and colorectal cancer survivorship care: perceptions of primary care physicians and medical oncologists in the United States. *J Clin Oncol* 31:2322-36, 2013
63. Mao JJ, Bowman MA, Stricker CT, et al: Delivery of Survivorship Care by Primary Care Physicians: The Perspective of Breast Cancer Patients. *Journal of Clinical Oncology* 27:933-938, 2009
64. Nyarko E, Metz JM, Nguyen GT, et al: Cancer survivors' perspectives on delivery of survivorship care by primary care physicians: an internet-based survey. *BMC Fam Pract* 16:143, 2015
65. Cheung WY, Aziz N, Noone AM, et al: Physician preferences and attitudes regarding different models of cancer survivorship care: a comparison of primary care providers and oncologists. *J Cancer Surviv* 7:343-54, 2013
66. Cheung WY, Neville BA, Cameron DB, et al: Comparisons of patient and physician expectations for cancer survivorship care. *J Clin Oncol* 27:2489-95, 2009

67. Shalom MM, Hahn EE, Casillas J, et al: Do survivorship care plans make a difference? A primary care provider perspective. *J Oncol Pract* 7:314-8, 2011
  68. Forsythe LP, Parry C, Alfano CM, et al: Use of survivorship care plans in the United States: associations with survivorship care. *J Natl Cancer Inst* 105:1579-87, 2013
  69. Chrischilles EA, McDowell BD, Rubenstein L, et al: Survivorship care planning and its influence on long-term patient-reported outcomes among colorectal and lung cancer survivors: the CanCORS disease-free survivor follow-up study. *J Cancer Surviv* 9:269-78, 2015
  70. Maly RC, Liang L-J, Liu Y, et al: Randomized Controlled Trial of Survivorship Care Plans Among Low-Income, Predominantly Latina Breast Cancer Survivors. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 35:1814-1821, 2017
  71. McDonough AL, Rabin J, Horick N, et al: Practice, Preferences, and Practical Tips From Primary Care Physicians to Improve the Care of Cancer Survivors. *Journal of Oncology Practice:JOP*.18.00740, 2019
  72. Reb A, Ruel N, Fakih M, et al: Empowering survivors after colorectal and lung cancer treatment: Pilot study of a Self-Management Survivorship Care Planning intervention. *Eur J Oncol Nurs* 29:125-134, 2017
  73. Mayer DK, Birken SA, Check DK, et al: Summing it up: An integrative review of studies of cancer survivorship care plans (2006-2013). *Cancer* 121:978-996, 2015
  74. Grunfeld E, Julian JA, Pond G, et al: Evaluating survivorship care plans: results of a randomized, clinical trial of patients with breast cancer. *J Clin Oncol* 29:4755-62, 2011
  75. Birken SA, Urquhart R, Munoz-Plaza C, et al: Survivorship care plans: are randomized controlled trials assessing outcomes that are relevant to stakeholders? *J Cancer Surviv* 12:495-508, 2018
  76. Selove R, Birken SA, Skolarus TA, et al: Using Implementation Science to Examine the Impact of Cancer Survivorship Care Plans. *Journal of Clinical Oncology* 34:3834-3837, 2016
  77. Parry C, Kent EE, Forsythe LP, et al: Can't see the forest for the care plan: a call to revisit the context of care planning. *J Clin Oncol* 31:2651-3, 2013
  78. Oeffinger KC, McCabe MS: Models for delivering survivorship care. *J Clin Oncol* 24:5117-24, 2006
  79. Nekhlyudov L, O'Malley D M, Hudson SV: Integrating primary care providers in the care of cancer survivors: gaps in evidence and future opportunities. *Lancet Oncol* 18:e30-e38, 2017
  80. Alfano CM, Mayer DK, Bhatia S, et al: Implementing personalized pathways for cancer follow-up care in the United States: Proceedings from an American Cancer Society-American Society of Clinical Oncology summit. *CA Cancer J Clin* 69:234-247, 2019
  81. National Health Service. (2012). London UK. Stratified Pathways of Care - From Concept to Innovation. <https://www.england.nhs.uk/improvement-hub/wp-content/uploads/sites/44/2017/11/Stratified-Pathways-of-Care.pdf>.
  82. National Health Service Improving Quality - London UK. (2013). Stratified Cancer Pathways: Redesigning Services for Those Living With or Beyond Cancer. <https://www.nice.org.uk/>.
  83. Jefford M, Rowland J, Grunfeld E, et al: Implementing improved post-treatment care for cancer survivors in England, with reflections from Australia, Canada and the USA. *Br J Cancer* 108:14-20, 2013
  84. Macmillan Cancer Support. Evaluation of the Transforming Cancer Follow-Up Programme in Northern Ireland. Final Report - London UK. <https://www.macmillan.org.uk/>.
- NIH Behavioral and Social Intervention Clinical Trial Protocol Template v3.0 - 20180827

85. Mead H, Pratt-Chapman M, Cleary S, et al: Evaluating the patient-centered quality of cancer survivorship care models. *Journal of Clinical Oncology* 35:48-48, 2017
86. Mead H, Pratt-Chapman M, Gianattasio K, et al: Identifying models of cancer survivorship care. *Journal of Clinical Oncology* 35:1-1, 2017
87. Campbell MK, Tessaro I, Gellin M, et al: Adult cancer survivorship care: experiences from the LIVESTRONG centers of excellence network. *J Cancer Surviv* 5:271-82, 2011
88. Shapiro CL, McCabe MS, Syrjala KL, et al: The LIVESTRONG Survivorship Center of Excellence Network. *J Cancer Surviv* 3:4-11, 2009
89. Cohen HJ: A model for the shared care of elderly patients with cancer. *J Am Geriatr Soc* 57 Suppl 2:S300-2, 2009
90. McCabe MS, Partridge AH, Grunfeld E, et al: Risk-based health care, the cancer survivor, the oncologist, and the primary care physician. *Semin Oncol* 40:804-12, 2013
91. Miller KD, Pandey M, Jain R, et al: Cancer Survivorship and Models of Survivorship Care: A Review. *Am J Clin Oncol* 38:627-33, 2015
92. O'Malley D, Hudson SV, Nekhlyudov L, et al: Learning the landscape: implementation challenges of primary care innovators around cancer survivorship care. *Journal of cancer survivorship : research and practice* 11:13-23, 2017
93. Overholser LS, Moss KM, Kilbourn K, et al: Development of a Primary Care-Based Clinic to Support Adults With a History of Childhood Cancer: The Tactic Clinic. *J Pediatr Nurs* 30:724-31, 2015
94. Cancer Survivorship E-Learning Series for Primary Care Providers; the National Cancer Survivorship Resource Center at George Washington University, in collaboration with the American Cancer Society: <https://smhs.gwu.edu/gwci/survivorship/ncsrc/elearning>. Accessed June 5, 2019.
95. Long-Term Survivorship Care After Cancer Treatment: Proceedings of a Workshop video, National Academy of Medicine <http://www.nationalacademies.org/hmd/Reports/2018/long-term-survivorship-care-after-cancer-treatment-proceedings.aspx>.
96. Cancer Patient to Cancer Survivor: Lost in Transition video , National Academy of Medicine <http://www.nationalacademies.org/hmd/Reports/2005/From-Cancer-Patient-to-Cancer-Survivor-Lost-in-Transition/From-Cancer-Patient-to-Cancer-Survivor-Lost-In-Transition.aspx>.
97. Cancer Survivorship in Primary Care, <http://www.cancerpcp.org/>, Accessed June 6, 2019.
98. American Society of Clinical Oncology Treatment Plan and Summary Resources: <https://www.asco.org/practice-guidelines/cancer-care-initiatives/prevention-survivorship/survivorship-compendium>, Accessed June 6, 2019.
99. Gould MK, Munoz-Plaza CE, Hahn EE, et al: Comorbidity Profiles and Their Effect on Treatment Selection and Survival among Patients with Lung Cancer. *Annals of the American Thoracic Society* 14:1571-1580, 2017
100. Hahn EE, Gould MK, Munoz-Plaza CE, et al: Understanding Comorbidity Profiles and Their Effect on Treatment and Survival in Patients With Colorectal Cancer. *J Natl Compr Canc Netw* 16:23-34, 2018



101. Hahn EE, Tang T, Lee JS, et al: Use of imaging for staging of early-stage breast cancer in two integrated health care systems: adherence with a choosing wisely recommendation. *J Oncol Pract* 11:e320-8, 2015
  102. Hahn EE, Wu YL, Munoz-Plaza CE, et al: Use of recommended posttreatment services for adolescent and young adult survivors of Hodgkin lymphoma. *Cancer* 125:1558-1567, 2019
  103. Chubak J, Yu O, Pocobelli G, et al: Administrative data algorithms to identify second breast cancer events following early-stage invasive breast cancer. *J Natl Cancer Inst* 104:931-40, 2012
  104. Hassett MJ RD, Taback N, Carroll N, Cronin AM, Ting GV, Schrag D, Warren JL, Hornbrook MC, Weeks JC: Validating billing/encounter codes as indicators of lung, colorectal, breast, and prostate cancer recurrence using two large contemporary cohorts. *Medical care research and review : MCRR* 52, 2014
  105. Dillman DA: *Mail and Internet surveys: The tailored design method*. . New York, NY, Wiley, 2007
  106. Yabroff KR, Dowling E, Rodriguez J, et al: The Medical Expenditure Panel Survey (MEPS) experiences with cancer survivorship supplement. *J Cancer Surviv* 6:407-19, 2012
  107. European Organisation for the Research and Treatment of Cancer Quality of Life Group, The EORTC QLQ-C30 Manuals, Reference Values and Bibliography, 2002 Brussels, Belgium.
  108. Hays RD, Martino S, Brown JA, et al: Evaluation of a Care Coordination Measure for the Consumer Assessment of Healthcare Providers and Systems (CAHPS) Medicare survey. *Medical care research and review : MCRR* 71:192-202, 2014
  109. Cella D, Riley W, Stone A, et al: The Patient-Reported Outcomes Measurement Information System (PROMIS) developed and tested its first wave of adult self-reported health outcome item banks: 2005-2008. *J Clin Epidemiol* 63:1179-94, 2010
  110. Blanch-Hartigan D, Chawla N, Beckjord EI, et al: Cancer survivors' receipt of treatment summaries and implications for patient-centered communication and quality of care. *Patient Educ Couns* 98:1274-9, 2015
  111. National Cancer Institute. Health Information National Trends Survey (HINTS): Survey instruments. <https://hints.cancer.gov/data/survey-instruments.aspx>.
  112. Pearman TP, Beaumont JL, Mroczek D, et al: Validity and usefulness of a single-item measure of patient-reported bother from side effects of cancer therapy. *Cancer* 124:991-997, 2018
  113. Holmes-Rovner M, Kroll J, Schmitt N, et al: Patient satisfaction with health care decisions: the satisfaction with decision scale. *Med Decis Making* 16:58-64, 1996
  114. Casillas J, Syrjala KL, Ganz PA, et al: How confident are young adult cancer survivors in managing their survivorship care? A report from the LIVESTRONG™ Survivorship Center of Excellence Network. *Journal of Cancer Survivorship* 5:371-381, 2011
  115. Schnipper LE, Lyman GH, Blayney DW, et al: American Society of Clinical Oncology 2013 top five list in oncology. *J Clin Oncol* 31:4362-70, 2013
  116. Schnipper LE, Smith TJ, Raghavan D, et al: American Society of Clinical Oncology identifies five key opportunities to improve care and reduce costs: the top five list for oncology. *J Clin Oncol* 30:1715-24, 2012
  117. Runowicz CD, Leach CR, Henry NL, et al: American Cancer Society/American Society of Clinical Oncology Breast Cancer Survivorship Care Guideline. *Journal of Clinical Oncology* 34:611-635, 2016
  118. Snyder CF, Frick KD, Peairs KS, et al: Comparing care for breast cancer survivors to non-cancer controls: a five-year longitudinal study. *Journal of general internal medicine* 24:469-474, 2009
- NIH Behavioral and Social Intervention Clinical Trial Protocol Template v3.0 - 20180827

119. Groenwold RHH, White IR, Donders ART, et al: Missing covariate data in clinical research: when and when not to use the missing-indicator method for analysis. CMAJ : Canadian Medical Association journal = journal de l'Association medicale canadienne 184:1265-1269, 2012